



Review

The microbial ecology of wine grape berries

A. Barata*, M. Malfeito-Ferreira, V. Loureiro

Laboratório de Microbiologia, Centro de Botânica Aplicada à Agricultura, Technical University of Lisbon, 1349-017 Lisbon, Portugal

ARTICLE INFO

Article history:

Received 8 July 2011

Received in revised form 28 October 2011

Accepted 27 November 2011

Available online 2 December 2011

Keywords:

Microbial ecology

Grapes

Damaged grapes

Spoilage yeasts

Acetic acid bacteria

Lactic acid bacteria

ABSTRACT

Grapes have a complex microbial ecology including filamentous fungi, yeasts and bacteria with different physiological characteristics and effects upon wine production. Some species are only found in grapes, such as parasitic fungi and environmental bacteria, while others have the ability to survive and grow in wines, constituting the wine microbial consortium. This consortium covers yeast species, lactic acid bacteria and acetic acid bacteria. The proportion of these microorganisms depends on the grape ripening stage and on the availability of nutrients. Grape berries are susceptible to fungal parasites until *véraison* after which the microbiota of truly intact berries is similar to that of plant leaves, which is dominated by basidiomycetous yeasts (e.g. *Cryptococcus* spp., *Rhodotorula* spp., *Sporobolomyces* spp.) and the yeast-like fungus *Aureobasidium pullulans*. The cuticle of visually intact berries may bear microfissures and softens with ripening, increasing nutrient availability and explaining the possible dominance by the oxidative or weakly fermentative ascomycetous populations (e.g. *Candida* spp., *Hanseniaspora* spp., *Metschnikowia* spp., *Pichia* spp.) approaching harvest time. When grape skin is clearly damaged, the availability of high sugar concentrations on the berry surface favours the increase of ascomycetes with higher fermentative activity like *Pichia* spp. and *Zygoascus hellenicus*, including dangerous wine spoilage yeasts (e.g. *Zygosaccharomyces* spp., *Torulaspota* spp.), and of acetic acid bacteria (e.g. *Gluconobacter* spp., *Acetobacter* spp.). The sugar fermenting species *Saccharomyces cerevisiae* is rarely found on unblemished berries, being favoured by grape damage. Lactic acid bacteria are minor partners of grape microbiota and while being the typical agent of malolactic fermentation, *Oenococcus oeni* has been seldom isolated from grapes in the vineyard. Environmental ubiquitous bacteria of the genus *Enterobacter* spp., *Enterococcus* spp., *Bacillus* spp., *Burkholderia* spp., *Serratia* spp., *Staphylococcus* spp., among others, have been isolated from grapes but do not have the ability to grow in wines. Saprophytic moulds, like *Botrytis cinerea*, causing grey rot, or *Aspergillus* spp., possibly producing ochratoxin, are only active in the vineyard, although their metabolites may affect wine quality during grape processing.

The impact of damaged grapes in yeast ecology has been underestimated mostly because of inaccurate grape sampling. Injured berries hidden in apparently sound bunches explain the recovery of a higher number of species when whole bunches are picked. Grape health status is the main factor affecting the microbial ecology of grapes, increasing both microbial numbers and species diversity. Therefore, the influence of abiotic (e.g. climate, rain, hail), biotic (e.g. insects, birds, phytopathogenic and saprophytic moulds) and viticultural (e.g. fungicides) factors is dependent on their primary damaging effect.

© 2011 Elsevier B.V. All rights reserved.

Contents

1.	Introduction	244
1.1.	Technological significance of microbial grape contaminants	244
2.	Methods to recover and identify grape microbial species	245
2.1.	Sampling schemes	245
2.2.	Sample picking and treatment	245
2.3.	Enrichment cultures	245
2.4.	Culture media	247
2.5.	Incubation conditions and colony selection	247
2.6.	Identification methods	247
2.6.1.	Molecular methods for yeast species identification	247
2.6.2.	Molecular methods for LAB species identification	248

* Corresponding author.

E-mail address: abbarata@isa.utl.pt (A. Barata).

2.6.3.	Molecular techniques for the rapid identification of AAB	248
2.6.4.	Other molecular and instrumental techniques	248
3.	Characterisation of grape microbial populations	248
3.1.	Population quantification	248
3.2.	Species diversity	248
3.3.	The case of <i>Saccharomyces</i> spp. and wine spoilage yeasts	252
3.4.	Habitat shared by other microbial species	252
4.	Factors influencing species diversity	252
4.1.	Climatic conditions	252
4.2.	Vineyard treatments	253
4.3.	Biotic factors	253
4.3.1.	Microbial vectors	253
4.3.2.	Microbial interactions and enzymatic activities	253
4.4.	Geographic location and vineyard factors: age, size, grape variety and vintage year	254
5.	Influence of grape damage or overripeness on grape juice and wine quality	254
5.1.	Dried or overripe grapes	254
5.2.	Noble-rot grapes	255
5.3.	Grey rot	255
5.4.	Powdery mildew	255
5.5.	Diffuse powdery mildew (DPM)	255
5.6.	Sour rot	255
5.7.	Other forms of damages	256
6.	Future prospects	256
	Acknowledgements	256
	References	256

1. Introduction

Grapes used for making wine have been studied for several decades. The earlier work of Louis de Pasteur, by the last quarter of the XIXth century, already showed that the microorganisms responsible for wine fermentations are yeasts present on the grapes. Since then a huge amount of information has been gathered on yeast dissemination in wine associated environments but ecological relationships are still to be fully understood. The prime focus has been the study of *Saccharomyces cerevisiae* given its relevance to winemaking. Earlier isolations using grape juice fermentations lead to the conclusion that this species was frequent in grapes (as reviewed by Vaughan-Martini and Martini, 1995). However, the use of direct isolation techniques showed a much different picture, revealing that *Saccharomyces* spp. is an absent or rare contaminant of grapes (Davenport, 1973, 1974). This fact has led to a still lively debate on the origin of the yeasts responsible for wine fermentation, being hypothesised that *S. cerevisiae* may be regarded as the first domesticated microbe (Martini, 1993).

Pioneering works on grape microbial ecology describe the main yeast species isolated from grapes and their environs (as reviewed by Fleet et al., 2002), while bacteria have been less studied. The recent increase in the number of grape-related yeast ecological studies did not contribute to a significant body of new knowledge. On the other hand, other areas of yeast ecology, like those related with plants, insects, soil or extreme habitats, have witnessed a significant development (Rosa and Péter, 2006) which provide useful guidance to study grape microbiota. In fact, most recent grape-related works are mostly dissemination studies. Microbial ecology is a wider concept, studying the ecosystems with their microbial interactions, microbial vectors, sources and sinks of microorganisms (Boddy and Wimpenny, 1992). Herman Phaff, the pioneer of yeast ecology, described ecology as “where microbes live and why they live in one habitat and how yeasts interact with other microorganisms” (Lachance, 2003). This global approach has been scarcely adopted to study the vineyard environment (Fleet et al., 2002). There are large gaps in the knowledge of the diversity and significance of yeast–yeast, yeast–fungi and yeast–bacteria interactions on grapes (Fleet, 2003). This may be because the microorganisms relevant to vine

and grapes diseases (phytopathogenic parasitic and saprophytic fungi) do not survive and grow in wine, while the species of the wine microbial consortium (WMC: yeasts, acetic acid bacteria and lactic acid bacteria) are not responsible for those diseases. The apparent rising of sour rot incidence, where the agents are part of the WMC (Barata et al., 2008a, in press), justify a reappraisal of the microbial ecology of grapes.

In this review we will attempt to make an update of the current awareness on the microbial ecology of grape surfaces, describing the species diversity, the factors affecting the species balance and the interactions with other microorganisms and vectors. We will also evaluate the current methods for microbial recovery and identification, trying to explain the contradictory results often found in literature. Special attention will be paid to the distribution of WMC species responsible for wine spoilage and to the effect of grape diseases on wine quality, given their technological importance. The question of the origin and dissemination of the WMC in the environment will be discussed aiming to provide future research directions.

1.1. Technological significance of microbial grape contaminants

The microbial species recovered from grapes may be divided into several groups according to their technological significance in grape and wine production. Under this view, microorganisms are characterised as a function of their effect on grape and wine quality.

The vine plant and grapes may be affected by a series of diseases of which the most well known are downy mildew (*Plasmopara viticola*), powdery mildew (*Erysiphe necator*) and grey rot (*Botrytis cinerea*), which are mostly prevented by phytochemical application. In addition, grapes may also bear saprophytic moulds (e.g. *Cladosporium* spp., *Aspergillus* spp., *Penicillium* spp.) responsible for several grape rots or mycotoxin production. However, these fungi do not have the ability to grow in wines and their effect on wine quality is due to grape damage. Contrarily, the microorganisms of the WMC are able to survive or grow on wine, depending on the efficiency of adequate processing measures. Thus, based on the concept already suggested by the authors (Loureiro and Malfeito-Ferreira, 2003; Malfeito-Ferreira, 2011), the species in this consortium may be grouped into: (i) easily controllable or innocent species, without the ability to

spoil wine when good manufacturing practises (GMP's) are applied; (ii) fermenting species, responsible for sugar and malic acid conversion; and (iii) spoilage *sensu stricto* species responsible for wine alteration even when GMP's are believed to be applied. Table 1 lists the most relevant microorganisms of referred groups, including recognised empirical denominations.

Concerning yeasts, the innocent group includes basidiomycetous species which are regarded as irrelevant to winemaking due to their inability to ferment juice sugars or to survive in wines. The ascomycetous dimorphic fungus, *Aureobasidium pullulans* (also called black yeast), a common yeast-like species, is also technologically irrelevant. The oxidative, weakly fermentative or fermentative ascomycetous species (*Candida* spp., *Kloeckera apiculata/Hanseniaspora uvarum*, *Metschnikowia* spp., *Pichia* spp.) are present in pre-fermentation steps or at the beginning of fermentation. The fermentation ability is not a well defined taxonomic feature and several species may be regarded as weakly fermentative or not. Among these, apiculate yeasts are determined by their microscopical shape and some strains may produce off-flavours in juices before or during fermentation (Romano, 2002). Film-forming yeasts (e. g. *Pichia*) owe their denomination to the ability of forming pellicles on the surface of bulk wines, being common contaminants of grapes, juices and wines, with the ability to produce off-flavours. Both apiculate and film-forming yeasts are regarded as contaminants because good manufacturing practises prevent their activity. Fermentative yeasts include those responsible for wine fermentation, where *S. cerevisiae* is the most important, but other species (*S. bayanus*, *S. pastorianus* and *S. paradoxus*) may also conduct or participate in the process (Skelin et al., 2008; Arroyo-López et al., 2010). These species may also be seen as wine spoilers if their activity persists beyond the fermentative steps of wine or sparkling wine production. The exception is that of *Saccharomyces* spp. forming desirable films on the surface of particular oxidative ageing processes of Sherry-like wines (Farris et al., 2002). Additionally, fermentative species comprise the spoilage *sensu stricto* yeasts which are technologically relevant due to their ability to spoil wines, either by off-flavour production (e. g. *Dekkera bruxellensis*) or sediment and cloudiness formation (e. g. *Zygosaccharomyces bailii*), under conditions following the GMP rules.

Concerning bacterial species, acetic acid bacteria are regarded as innocent because they are easily controllable by GMP's in the winery, although the exaggerated production of acetic acid during grape sour rot is a serious threat to wine quality. The physiological diversity of lactic acid bacteria does not allow a precise assessment of their technological significance. The typical agent of malolactic fermentation is *O. oeni* while *Lactobacillus* spp. and *Pediococcus* spp. may be responsible for spontaneous fermentations (Lerm et al., 2010). These species may spoil wine when their activity goes beyond malic conversion, particularly in high pH wines, producing off-flavours or biogenic amines (Arena et al., 2011; Capozzi et al., 2011; Pan et al., 2011). Grapes also bear a wide diversity of bacterial species common in nature or in other food related environments (Table 1). However, they should not be integrated in the WMC because they do not influence wine quality.

2. Methods to recover and identify grape microbial species

Several reviews on the analytical approaches to study overall yeast ecology have been published elsewhere (Boundy-Mills, 2006; Ciani et al., 2002; Kurtzman et al., 2011). We will try to extend the discussion to bacterial species and to the issues related to grape analysis, taking in consideration the advice of Lachance (2003) to obtain ecologically meaningful conclusions: (i) adequate sample size, (ii) correct identification, (iii) habitat characterisation, (iv) substrate sampling, that is far more important than the sampling procedure, and (v) sample replication, which is more important than serial dilutions and plate counts.

2.1. Sampling schemes

Yeast populations in nature suffer spatial and temporal fluctuations (Fonseca and Inácio, 2006) that must be taken into account when devising sampling schemes. Ecological conclusions must be based on extensive sampling. In fact, different bunches of grapes bear different populations and it is more informative to analyse several smaller samples than to blend several bunches into a larger one (Barata et al., 2008b). Samples should be taken in several locations in the vineyard so that spatial fluctuations dictated by the uneven microbial distribution are minimised. Repeated sampling over the years is also a sound practise to understand the behaviour of natural microbial populations and avoid unsupported conclusions.

2.2. Sample picking and treatment

Once sampling schemes are defined, the first step in grape analysis concerns the choice of the method to pick the grapes. The main concern should be to aseptically separate sound berries from damaged ones given their completely different microbial load. This cannot be achieved if whole bunches are analysed because damaged berries may be hidden inside. As an example, a sound bunch containing 50 sound berries with an average of 10^3 CFU/berry would contain 2×10^4 CFU/berry if only one damaged berry with 10^6 CFU/berry blended together with the sound berries. Further, the number of species of the "sound" bunches would reflect the diversity of the damaged berry and not of the whole sound bunch. Berries may be picked separately in the vineyard, or grape bunches are collected in the field and the berries are separated in the laboratory to give a 300 g sample. When bunches are picked and taken to the laboratory, sound berries should be chosen from fully sound bunches after visual inspection because sound berries from partially damaged bunches bear higher yeast numbers (Barata et al., 2008a).

The isolation from grape juices obtained in the winery (industrial or experimental) may only be regarded as an approximation of the natural grape microbiota. In fact, when grape juice is only sampled in the winery, even at the beginning of fermentation, a different picture of yeasts species may be obtained, due to bulk transport, crushing or pressing in the winery (Barata et al., in press; Sturm et al., 2006). Therefore, it is mandatory to isolate yeasts from grapes aseptically collected in the vineyard and they must be processed as quickly as possible because grapes damaged during transport to laboratory may accumulate higher numbers of yeasts (Yanagida et al., 1992).

The method for yeast dislodgement from grapes is also of importance. Belin (1972) discussed the efficiency of recovery methods, saying that simple washings with shaking may not be enough. Later, Martini et al. (1980) presented evidence of the importance of strong disruptive methods followed by enrichment cultures to obtain an exhaustive picture of the yeast flora. Direct enrichment gave lower number of species, than washing and sonication, but enrichment may be the only way to recover fermenting species. These authors advised enrichment in one sample, and agitation or percolation followed by sonication in another sample, but they did not try grape blending. This method was performed by Combina et al. (2005) who found higher results with grape blending in plastic bags, than with jet streaming and shaking. Prakitchaiwattana et al. (2004) evaluated recovery during 4 successive rinsings for 10 min each. Rinsing released 80% of total yeasts in the first wash and 96% in damaged grapes. Differences in species diversity through all steps were not apparent or were due to a dilution effect. Therefore, the classical food sample suspension in peptone or saline solutions followed by stomaching and serially diluting is a reliable option.

2.3. Enrichment cultures

When juice is obtained after grape blending or when suspensions are recovered after grape or single berry washing, the following step

Table 1
Dissemination and technological significance of microbial species isolated from the vineyard and winery environment.
Data compiled from Loureiro and Malfeito-Ferreira, 2003; Malfeito-Ferreira, 2011, and Loureiro et al., in press).

Group	Metabolism	Genus	Relevant species	Technological significance	Main Sources	Occasional sources
Filamentous fungi	Obligate parasites		<i>Plasmodium viticola</i> <i>Erysiphe necator</i>	Downy mildew Powdery mildew	Vine tissues Vine tissues	
	Saprophytic moulds	<i>Botrytis</i> <i>Aspergillus</i>	<i>B. cinerea</i> <i>A. alliaceus</i> , <i>A. carbonarius</i> , <i>A. niger</i> <i>aggregate</i> , <i>A. ochraceus</i>	Grey rot, Noble rot <i>Aspergillus</i> rot, ochratoxin A producers	Ubiquitous Ubiquitous	
		<i>Penicillium</i>	<i>P. expansum</i>	Green mould, patulin producer	Ubiquitous	
		<i>Cladosporium</i> <i>Colletotrichum</i> <i>Greeneria</i>	<i>C. herbarum</i> <i>C. acutatum</i> <i>G. uvicola</i>	<i>Cladosporium</i> rot Ripe rot Bitter rot	Ubiquitous Ubiquitous Ubiquitous	
Yeasts						
Basidiomycetous	Oxidative	<i>Filobasidium</i> , <i>Cryptococcus</i> , <i>Rhodotorula</i> (pink yeast)		Absent/unknown	Soil, bark, leaf, grape	Grape juice
Ascomycetous	Oxidative	<i>Aureobasidium</i>	<i>A. pullulans</i> (yeast-like fungi, black yeast)	Absent/unknown	Soil, bark, leaf, grape	Grape juice
	Oxidative or weakly fermentative	<i>Hanseniaspora/Kloeckera</i> (apiculate yeast)	<i>H. uvarum</i> / <i>K. apiculata</i>	Contamination/spoilage	Grape, grape juice, fermentation	Soil
		<i>Candida</i> (film-forming yeast)	<i>C. stellata</i> or <i>C. zemplinina</i>	Contamination	Grape, grape juice, fermentation, wine	Bark, soil, insects
	Fermentative	<i>Metschnikowia</i> <i>Pichia</i> (film-forming yeast)	<i>Zygoascus hellenicus</i> / <i>C. steatolytica</i>	Contamination	Grape, grape juices	Fermentation, insects
		<i>Debaryomyces</i> <i>Lachancea</i> (ex <i>Kluyveromyces</i>)	<i>M. pulcherrima</i> <i>P. anomala</i> <i>P. membranifaciens</i>	Contamination Contamination/spoilage Contamination/spoilage	Grape, fermentation Grape, fermentation Grape, fermentation, wine	Insects
		<i>Torulaspota</i>	<i>P. guilliermondii</i>	Contamination/spoilage	Grape, fermentation, wine	
		<i>Zygosaccharomyces</i>	<i>D. hansenii</i> <i>L. thermotolerans</i> <i>L. fermentati</i> (ex <i>Z. fermentati</i>) <i>T. delbrueckii</i>	Contamination Contamination	Grape, fermentation Grape, fermentation	
		<i>Zygosaccharomyces</i>	<i>Z. bailii</i>	Spoilage	Wine, concentrated grape juices	Grape, fermentation
		<i>Dekkera/Brettanomyces</i> <i>Saccharomyces</i>	<i>Z. bisporus</i>	Spoilage	Wine	Grape, fermentation
		<i>Schizosaccharomyces</i>	<i>Z. rouxii</i>	Spoilage	Concentrated grape juices	
		<i>Saccharomyces</i>	<i>D. bruxellensis</i> <i>S. cerevisiae</i> <i>S. bayanus</i> <i>S. paradoxus</i> <i>S. pastorianus</i> <i>Sc. pombe</i>	Spoilage Fermenting/spoilage Fermenting/spoilage	Wine Fermentation, wine Fermentation, wine Fermentation, wine Fermentation, wine	Grape, insects Soil, Grape Grape Grape Grape
		<i>Saccharomyces</i>	<i>S. ludwigii</i>	Spoilage	Wine	Grape, fermentation Grape, fermentation
Bacteria						
Acetic acid bacteria	Aerobic	<i>Gluconobacter</i> spp., <i>Acetobacter</i> spp., <i>Gluconoacetobacter</i> spp.,		Wine spoilage, vinegar production	Grape, wine	Insects
Lactic acid bacteria	Anaerobic, semi-anaerobic	<i>Oenococcus</i> sp., <i>Lactobacillus</i> spp., <i>Pediococcus</i> spp., <i>Weissella</i> spp.		Malolactic fermentation or wine spoilage	Grape, wine	
Several bacterial species		<i>Acinetobacter</i> spp., <i>Curtobacterium</i> spp., <i>Pseudomonas</i> spp., <i>Serratia</i> spp., <i>Enterobacter</i> spp., <i>Enterococcus</i> spp., <i>Bacillus</i> spp., <i>Staphylococcus</i> spp.		Innocuous contaminants	Grapes	

is plating or to continue using enrichment cultures. If juice is allowed to ferment spontaneously this corresponds to an auto-enrichment step. True spontaneous fermentations are performed in the absence of sulphur dioxide additions. If this is added, sensitive species are affected. The enrichment may also be done with culture media where single berries, diluted suspensions or juices are introduced.

The enrichment step elicits the recovery of minority species which would not be detected by plating. The typical example is the wine fermenting *S. cerevisiae*, as mentioned before, but the recovery of other technologically relevant species also requires this approach. Renouf and Lonvaud-Funel (2007) only detected the dangerous spoilage yeast *D. bruxellensis* by using a selective medium as an enrichment step. Washing solutions of grapes did not reveal this species. The enrichment step also enabled the detection of *S. cerevisiae* and other fermentative species (*C. cantarelli* and *P. fermentans*) while oxidative basidiomycetous species (*Cryptococcus laurentii*, *Rhodotorula mucilaginosa* and *Sporidiobolus pararoseus*) were detected by these authors in the washing solution. The isolation of the malolactic agent *O. oeni* also required this type of approach using washing solutions (Renouf et al., 2005) or enrichment broths (Renouf et al., 2007). The matter of appropriate culture media may be required to isolate *O. oeni* but using enrichment broths Bae et al. (2006) did not manage to recover this species. On the contrary, these authors detected species of *Lactobacillus*, *Enterococcus*, *Lactococcus* and *Weissella* using enrichments broths.

2.4. Culture media

The choice of culture media is critical for characterising complex microbial populations (Loureiro et al., 2004). The utilisation of general purpose culture media directly from grape juice or dilutions only enables the recovery of the most frequent and faster growing species. Spread plates using a 0.1 ml sample are commonly preferred and colonies of the most representative species occupy the medium surface and cover minority colonies, leaving undetectable those representing less than about 1% of the population (Fleet et al., 2002).

The recovery of minority species and/or slow growers requires the use of selective media, possibly following sample enrichment. If the enrichment step is done with selective media then a general purpose medium may be used to isolate growing species because the overall populations were already restricted. Wine spoilage species are typically minority species and so it is advised to use selective media. Several reports mention selective media for *Schizosaccharomyces pombe* (Florenzano et al., 1977), *Z. bailii* (Schuller et al., 2000), *D. bruxellensis* (Rodrigues et al., 2001) and *S. cerevisiae* (Kish et al., 1983). Lactic acid bacteria are also minor constituents of berry microbiota and profit from the utilisation of specific media supplemented with ethanol or tomato juice (Bae et al., 2006).

The use of antifungal compounds is advisable to restrict the growth of filamentous fungi, although some yeast species may be affected (Loureiro et al., 2004). Further, the selection of adequate antibiotics is essential to detect separately yeasts, lactic acid and acetic acid bacteria. When moulds or environmental bacteria are numerous and a quantification of WMC species is wanted it is advisable to use the Most Probable Number (MPN) technique (Loureiro et al., 2004). With appropriate choice of culture media, antibiotics and incubation conditions, it is possible to inhibit, at least partially, the background microbiota and obtain numbers of the selected species (Barata et al., in press).

2.5. Incubation conditions and colony selection

After surface inoculation plates should be kept upright when yeasts are to be recovered, as opposed to bacteria (Deák and Beuchat, 1996). Incubation temperature is usually set at 25–30 °C, but lower temperatures (10 °C) are essential to recover *S. uvarum*

(Sampaio and Gonçalves, 2008). Aerobic conditions are preferred but anaerobiosis was advised for *Sc. pombe* (Florenzano et al., 1977) and lactic acid bacteria (Barata et al., in press). In liquid media, the availability of oxygen may also be varied by using different shaking rates. Another important issue is to use long incubation periods, up to 14 days, specially when wine spoilage yeasts are screened (Florenzano et al., 1977; Rodrigues et al., 2001; Schuller et al., 2000).

After incubation, strain choice by colony observation must be done by experienced technicians, small differences are frequently overlooked, being also indispensable to perform microscopical examinations. Frequently, same colony morphologies belong to different species, being a good practise to choose more than one colony for each morphological type.

2.6. Identification methods

Accurate identification is obviously crucial for the establishment of microbial communities. It is accepted that classical identification techniques based on morphological, biochemical and physiological criteria (Barnett et al., 1990; Kreger-van Rij, 1984; Krieg and Holt, 1984–1989) may have provided incorrect results in the past or could not reach species definition due to heterogeneous phenotypical results. This is particularly true for species of the genus poorly defined by conventional methodologies (e. g. *Candida* spp., *Pichia* spp.), and so it is not surprising that the number of their species is increasing in recent surveys. Moreover, the reproducibility of these techniques is somewhat questionable, since in many cases they depend on the physiological state of the cells. Molecular biological techniques circumvent these difficulties by allowing direct analysis of the genome, irrespective of the physiological state of the cell, providing more precise identifications.

2.6.1. Molecular methods for yeast species identification

The principles underlying yeast identification by molecular techniques have been previously addressed (Giudici and Pulvirenti, 2002). This theme is subjected to a permanent evolution and more recent reviews provide an adequate update of the available methodologies (Cocolin et al., 2011a; Fernández-Espinar et al., 2011; Querol and Fleet, 2006). The main technical alternatives for species identification are briefly described below, while approaches required for accurate studies of source tracking or evolutionary assessments are available elsewhere (Fernández-Espinar et al., 2011).

2.6.1.1. Sequencing of ribosomal DNA. Yeast species can be identified by comparison of nucleotide sequences from rDNA regions. The two most commonly used regions are the D1 and D2 regions encoding the 26S (Kurtzman and Robnett, 1998) and 18S (James et al., 1997) ribosomal subunits. The availability of sequences in DNA databases, particularly for the D1/D2 region of the 26S gene, makes this technique useful for assigning unknown yeast to a specific species when the homology of the sequence is greater than 99% (Kurtzman and Robnett, 1998).

2.6.1.2. Restriction analysis of ribosomal DNA (rDNA). Simpler methods have been designed based on PCR amplification of rDNA regions followed by restriction analysis of the amplified products. A very useful rDNA region that can be used to differentiate between species is that containing the 5.8S gene and the adjacent intergenic regions ITS1 and ITS2. This technique was used by Guillamón et al. (1998) for the rapid identification of wine yeasts, and was later extended to yeasts associated with foodstuffs and beverages (de Llanos et al., 2004; Esteve-Zarzoso et al., 1999; Fernández-Espinar et al., 2000). The amplified fragments and restriction profiles for these species with *HaeIII*, *HinfI*, *CfoI* and *DdeI* are available online at <http://yeast-id.com/>. In the absence of restriction profiles the solution is to sequence the ribosomal

DNA as described before. When restriction profiles are coincident, identification may also rely on the utilisation of classical biochemical tests (Barata et al., 2008a, 2008b).

2.6.1.3. Polymerase chain reaction (PCR)-denaturing gradient gel electrophoresis (DGGE). This genetic fingerprinting technique based on PCR amplification and denaturing gradient gel electrophoresis (DGGE) has been introduced into microbial ecology by Muyzer et al. (1993). This technique allows DNA fragments of the same length to be separated on the basis of sequence differences. DNA migration is retarded when the DNA strands dissociate at a specific concentration of denaturing agent. A related technique is temperature gradient gel electrophoresis (TGGE), which is based on a linear temperature gradient for separation of DNA molecules (Fernández-Espinar et al., 2011). The DGGE and TGGE methods have only recently been used, for yeast identification in wine fermentations (Andorrà et al., 2008; Cocolin et al., 2000; Di Maro et al., 2007; Prakitchaiwattana et al., 2004; Renouf et al., 2007; Urso et al., 2008). These techniques are directly applied to the sample but their low sensitivity is a major drawback to study minority populations.

2.6.2. Molecular methods for LAB species identification

A wide variety of molecular techniques have been used to characterise LAB from wine. These methods were recently reviewed by Munõz et al. (2011). Amplified rDNA restriction analysis (ARDRA) has been used as a rapid, reliable method of identifying the main LAB involved in winemaking (Rodas et al., 2003; Ventura et al., 2000).

Amplified rDNA fragment analysis via PCR followed by DGGE has also been used to compare diversity and monitor changes in populations of LAB during the winemaking process (Lopez et al., 2003). More recently, Rodas et al. (2005) concluded that the randomly amplified polymorphic DNA (RAPD) and ribotyping are useful for identifying and classifying LAB, whereas ARDRA is useful only for identification purposes and the restriction fragment length polymorphism-pulsed field gel electrophoresis (RFLP-PFGE) is useful for distinguishing between different LAB strains of the same species (Rodas et al., 2003).

2.6.3. Molecular techniques for the rapid identification of AAB

Sequencing is one of the methods proposed for the identification of AAB and generally involves ribosomal genes or the region between the 16S and 23S ribosomal genes (Yamada and Yukphan, 2008). The latter involves intergenic regions known as internal transcribed spacers (ITSs). New approaches were recently developed to detect and group different microorganisms using simple, rapid, and inexpensive techniques in order to reduce the cost of DNA sequencing in descriptive ecological studies evolving hundreds of samples (Guillamón and Mas, 2011).

Several molecular techniques have been developed for the rapid and reliable identification of most AAB species, particularly those present in grapes, wine, and vinegars. The first method involves restriction analysis of the 16S ribosomal gene following amplification by PCR using a protocol known as 16S-ARDRA (Poblet et al., 2000; Ruiz et al., 2000). More recently González et al. (2006a) proposed a method based on the RFLP-PCR of the 16S rRNA and 16S-23S ITS regions, involving the sequential use of different restriction enzymes to allow the grouping and distinguishing of the currently described AAB species. This approach has been used for the identification of AAB present in grapes (Prieto et al., 2007), wines (González et al., 2004, 2005) and vinegars (Gullo et al., 2006; Ilabaca et al., 2008).

Real-time quantitative PCR methods circumvents the problem of culturing AAB, which affects the previous culture depending techniques and have been successfully used for the identification and quantification of AAB in grapes, wine and vinegars (González et al., 2006b; Torija et al., 2010).

Finally, DGGE and TGGE can also be used for the identification of AAB in combination with PCR to separate amplified fragments of the

16S gene according to small sequence differences (Guillamón and Mas, 2011). DGGE (Andorrà et al., 2008; De Vero et al., 2006; Gullo et al., 2009; Lopez et al., 2003) and TGGE (Ilabaca et al., 2008) methods also circumvent the problem of culturing AAB and have been successfully used for the identification AAB in grapes, wine and vinegars.

2.6.4. Other molecular and instrumental techniques

Metagenomics is an increasingly used alternative, which does not require cloning or PCR amplification, and can produce huge numbers of DNA readings to study uncultured organisms (Huson et al., 2007). Its application to grape musts evidenced the expected microbiota diversity and the unexpected presence of some species unusual in wines, such as *Enterobacter* spp. (Nisiotou et al., 2011) or the green algae *Dunaliella tertiolecta* (Gomes et al., 2009).

Fourier transformed infrared spectroscopy has been reported as a powerful technique to identify yeasts, moulds and bacteria (Wenning et al., 2002, 2010; Santos et al., 2010) but we are not aware of reports concerning grape berry microbiota.

In conclusion, according to our experience, grape sampling is the critical step to obtain sound ecological data. The evaluation of the factors affecting the microbial ecology of grapes depends on berry selection according to its health status and on adequate experiment repetition during more than 2 annual cycles. In addition, species diversity is enhanced by the utilisation of selective media and enrichment techniques, particularly to recover minor populations. Identification can be achieved by molecular techniques chosen according to the targeted species. At present, direct molecular identification is only suitable for the dominant populations, being difficult to apply when samples are contaminated by a wide variety of moulds, yeasts and bacteria.

3. Characterisation of grape microbial populations

3.1. Population quantification

The yeast populations of grapes are roughly comprised between 10^2 and 10^4 cells/g (Fleet et al., 2002), but higher values have also been reported (Table 2). This wide range of values may be explained, at least partially, by bunch sampling without accurate separation of damaged berries, as explained before. When this is done, smaller variations are found (Barata et al., 2008a). Table 3 shows a range of total counts provided by surveys where samples were separated according to health status. As a rule, damage induced at least an increase of one log cycle. The availability of high sugar concentrations explains the higher populations on damaged grape berries. The exception is related to damage by honeydew produced by mealybugs, where counts are similar in sound and affected berries, probably because of antimicrobial compounds and high sugar concentration, as in honey produced by bees (Barata et al., 2008b).

Bacterial populations are usually several orders of magnitude lower than those of yeasts in sound grapes. Lactic acid bacteria have counts lower than 10^2 CFU/g (Barata et al., in press; Francesca et al., 2011), explaining the use of enrichment cultures for their current detection (Bae et al., 2006; Renouf et al., 2007), which do not seem to increase significantly on damaged grapes (Barata et al., in press).

Contrarily, acetic acid bacteria are stimulated by berry damage, increasing from less than 10 CFU/g to 10^6 CFU/g on rotten grapes (Barata et al., in press; Barbe et al., 2001). On the contrary, Nisiotou et al. (2011) found comparable acetic acid bacterial counts in sound and grey rotten grapes, ranging from 10^5 to about 10^6 CFU/mL.

3.2. Species diversity

Worldwide surveys seem to indicate that apparently sound grapes are colonised by a wide variety of yeast species without any obvious

Table 2

Yeast and yeast-like species isolated from “sound” grapes or berries at harvest (data collected from selected surveys, the symbol + indicates relative proportion of the detected species).

Species	France	France	Italy	Spain	Spain	Portugal	Greece	Slovenia	Canada	Brazil	Argentina	Japan	Australia	India	China
Basidiomycetes ^a	+++	+++	++	+++		+++		+++	+++	+	+	+++	++		+++
<i>Aureobasidium pullulans</i>	+		+++	+++	+		+	++	+++	+			+++		
<i>Hanseniaspora</i> spp.		++			+++		++					+		+++	+
<i>H. uvarum</i>	+++		++	+	+++	++	+++	++	++	+++	+++	+++		+	+++
<i>Metschnikowia</i> spp.	+		++		+	+		+			+++				+
<i>Candida</i> spp.		+++	+	++		+					+	++		++	++
<i>C. stellata/zemplinina</i>						+	++				+				+
<i>Debaryomyces</i> spp.		+						+						+	
<i>Issatchenkia</i> spp.		+			++		+			+++	+			++	+
<i>Kluyveromyces</i> spp./ <i>Lachancea</i> spp.		+			++										
<i>Pichia</i> spp.		+	+		+	+		+			++			++	++
<i>Brettanomyces</i> spp.			+												
<i>Saccharomyces</i> spp.		+	+												
<i>S. cerevisiae</i>					++									++	
<i>Saccharomycopsis</i> spp.			++												
<i>Saccharomycodes ludwigii</i>											+				
<i>Torulaspota</i> spp.					+	+									
<i>Zygosaccharomyces</i> spp.			+												+
<i>Z. bailii</i>					+										+
Sampling ^b	Be	Bu	Be	Be	Bu	Be	Bu	Bu	Bu	Be, Bu	Bu	Bu	Bu	Bu	Bu
Culture media ^c	G	G	G	G	G,A	G,S	G,S	G	G	G	G	–	G	G	G,S
Count range (log)	4–6/mL	3–5/berry	0.1–4/g	–	–	2–6/g	3–4/g	3–6	–	–	5–6/g	0.7–3/mL	2–4/g	–	3–5/g
References	Barnett et al. (1972)	Renouf et al. (2005)	Guerzoni and Marchetti (1987)	Sabate et al. (2002)	Clavijo et al. (2010)	Barata et al. (2008a, 2008b)	Nisiotou and Nychas (2007)	Raspor et al. (2006)	Subden et al. (2003)	Baffi et al. (2011)	Combina et al. (2005)	Yanagida et al. (1992)	Prakitichaiwattana et al. (2004)	Chavan et al. (2009)	Li et al. (2010)

^a *Cryptococcus* spp., *Bulleromyces* spp., *Sporidiobolus* spp., *Sporobolomyces* spp., *Rhodotorula* spp., *Trichosporon* spp.^b Be, berry, Bu, bunch.^c G, general purpose, A, autoenrichment, S, selective media.

Table 3
Yeast and yeast-like counts (range (x–y) or mean ± standard deviation (z ± t) of log CFU/g or log CFU/berry) and species frequency found in sound and damaged grapes at harvest.

Damage	Sound grapes	Species (frequency)	Damaged grapes	Species (frequency)	Reference
Undefined ^a	4.4–4.4 (2) ^b	<i>Aureobasidium pullulans</i> (95–100) <i>Cyptococcus victorae</i> (0–4) <i>Rhodotorula laryngis</i> (0–1)	5.9–7.2 (2) ^b	<i>Metschnikowia</i> spp. (64–75) <i>Hanseniaspora</i> spp. (0–36) <i>A. pullulans</i> (0–25)	Prakitichaiwattana (2004) ^c
Diffuse powdery mildew	2.3–4.0 (4)	<i>Aureobasidium</i> spp. (65–66) <i>Hanseniaspora</i> spp. (22–23)	4.0–6.0 (4)	<i>Hanseniaspora</i> spp. (67–80) <i>Metschnikowia</i> spp. (3–19) <i>Aureobasidium</i> spp. (2–5) <i>Candida stellata</i> (4–10)	Gadoury et al. (2007) ^c
Sour rot	3.2–3.9 (5)	<i>A. pullulans</i> (19) <i>Saccharomycopsis vini</i> (15) <i>Metschnikowia pulcherrima</i> (14) <i>Rhodotorula</i> (11) <i>H. uvarum</i> (11) <i>Pichia membranifaciens</i> (8) <i>Zygosaccharomyces</i> spp. (6) <i>C. krusei</i> (4)	6.6–6.8 (5)	<i>C. krusei</i> (33) ^a <i>Sac. vini</i> (19) <i>M. pulcherrima</i> (15) <i>Hanseniaspora uvarum</i> (11) <i>C. steatolytica</i> (10) <i>Zygosaccharomyces</i> spp. (3) <i>C. sorbosa</i> (3) <i>P. membranifaciens</i> (1) <i>A. pullulans</i> (<1)	Guerzoni and Marchetti (1987) ^d
	2.0–2.8 (2)	Basidiomycetes (0–100) <i>C. vanderwaltii</i> (0–11)	6.8–7.2 (2)	<i>C. vanderwaltii</i> (60–74) <i>H. guilliermondii</i> (3–24) <i>Lachancea thermotolerans</i> (2–6) <i>P. membranifaciens</i> (0.3–16) Basidiomycetes (0.1–0.3) <i>H. uvarum</i> (0–12) <i>Zygoascus hellenicus</i> (0 ≤ 0.1) <i>Candida</i> spp. (0–3) <i>Issatchenkia</i> spp. (0–1) <i>H. uvarum</i> (76) <i>C. zemplanina</i> (12) <i>Issatchenkia occidentalis</i> (5) <i>I. terricola</i> (3.5) <i>H. opuntiae</i> (3.5)	Barata et al. (2008a) ^c
Grey rot	4.3 ± 0.4 (3)	<i>H. uvarum</i> (67) <i>C. zemplanina</i> (22) <i>H. opuntiae</i> (6.5) <i>A. pullulans</i> (4.5)	5.7 ± 0.6 (3)	<i>H. uvarum</i> (75) <i>M. pulcherrima</i> (7) <i>H. opuntiae</i> (5.5) <i>H. guilliermondii</i> (3.5) <i>Z. bailii</i> (3.5) <i>C. zemplanina</i> (3.5) <i>I. terricola</i> (2)	Nisiotou and Nychas (2007) ^c
Noble rot	3.3 ± 0.7 (3)	<i>H. uvarum</i> (77) <i>H. guilliermondii</i> (9) <i>H. opuntiae</i> (9) <i>I. terricola</i> (5)	6.8 ± 0.8 (3)	<i>H. uvarum</i> (75) <i>M. pulcherrima</i> (7) <i>H. opuntiae</i> (5.5) <i>H. guilliermondii</i> (3.5) <i>Z. bailii</i> (3.5) <i>C. zemplanina</i> (3.5) <i>I. terricola</i> (2)	Nisiotou and Nychas (2007) ^c
Honeydew	3.7–5.9 (19)	Basidiomycetes (62) ^b <i>H. uvarum</i> (23) <i>M. pulcherrima</i> (6) <i>Candida</i> spp. (3) <i>C. stellata</i> (2) <i>C. vanderwaltii</i> (2) <i>Torulaspota pretoriensis</i> (2)	3.3–6.5 (26)	Basidiomycetes (68) ^b <i>Z. hellenicus</i> (10) <i>H. uvarum</i> (7) <i>C. vanderwaltii</i> (6) <i>Pichia</i> spp. (4) <i>T. pretoriensis</i> (2) <i>L. thermotolerans</i> (2) <i>M. pulcherrima</i> (1) <i>Sac. vini</i> (1) <i>Z. bisporus</i> (1)	Barata et al. (2008b) ^d

^a Analysis of intact berries or berries physically damaged, shrivelled, broken skin, mouldy (seldom) or discoloured.

^b Number of samples analysed given within brackets.

^c Frequency expressed as a percentage of specific colonies relative to total colony number on plate media.

^d Frequency expressed as a percentage of positive samples in relation to total samples analysed.

explanation (Table 2). However, this variety may be reduced to relatively few groups of similar physiological characteristics. For instance, the ubiquitous *Candida* spp. and *Pichia* spp. are highly heterogeneous, and new species are likely to be found in each new survey because the accuracy of molecular identifications is constantly increasing (Rao et al., 2007). In addition, these genera are ubiquitous and given the strong influence of the vineyard surroundings, it is also likely that a wide diversity of their species will be found. For instance, (Chavan et al. (2009) were the first to isolate *C. azyma* from grapes. This species was characteristic of the sugarcane formerly planted in the area of the studied vineyard which is likely to be the explanation for this outcome. Therefore, we propose to systematise the WMC microbiota of grape berries into three main yeast groups, characterised by similar behaviour on grape berries:

- (i) oligotrophic, oxidative basidiomycetous yeasts, the yeast-like fungi *A. pullulans*, and lactic acid bacteria (*Lactobacillus* spp., *Oenococcus oeni*);

- (ii) copiotrophic, oxidative ascomycetes (several *Candida* spp.); weakly fermentative apiculate (*Hanseniaspora* spp.), film-forming (*Pichia* spp.), fermentative (*C. zemplanina*, *Metschnikowia* spp.) yeasts;
- (iii) copiotrophic strongly fermentative yeasts (*Saccharomyces* spp., *Torulaspota* spp., *Zygosaccharomyces* spp., *Lachancea* spp. and *Pichia* spp.) and the obligate aerobic acetic acid bacteria (*Gluconobacter* spp., *Gluconoacetobacter* spp., *Acetobacter* spp.).

The balance among these groups, after *véraison*, is particularly dependent on nutrient availability on berry surface, as described below.

The first group is composed by species favoured by the nutrient poor environment of truly sound berries. The awareness that basidiomycetes are dominant on grape surfaces is relatively recent (Yanagida et al., 1992) and reflects the use of direct analysis of grapes before juice fermentation (Davenport, 1973, 1974). This predominance, similar to that of plant leaves (Fonseca and Inácio, 2006; Sabate et al., 2002), may be explained by their oligotrophy required

to survive in the harsh environment of the unblemished cuticle. These species are ubiquitous, spread all over the environment. In vineyards, soil, leaves and bark also characterised by the dominance of basidiomycetous oxidative yeasts and the yeast-like fungi *A. pullulans* (Davenport, 1974; Poulard et al., 1981; Sabate et al., 2002).

The increase in the proportion of oxidative or weakly fermentative ascomycetous species (*Hanseniaspora*, *Candida*, *Metschnikowia* and *Pichia* spp.), may occur during ripening. The mechanisms underlying this succession are not clear, species interaction may occur (Fleet, 2003), but the main factor should be related to nutrient availability. In fact, when approaching maturity, berries begin to behave differently from plant leaves, probably because of cuticle softening and release of volatile organic compounds (VOC's). The emergence of these species is probably the result of juice release, even in visually intact berries, as suggested by the effect of diffuse powdery mildew which injury is invisible to the naked eye (Table 3). This hypothesis has not been tested but the fact that mature sound berries may only harbour basidiomycetes and *A. pullulans* at harvest time (Barata et al., in press), suggests that some nutrient release should occur to enable growth of the copiotrophic biota. Moreover, we believe that many of the apparently contradictory results may be explained by using grape bunches without separating damaged berries. This has been demonstrated recently by Čadež et al. (2010). As expected, these authors found that basidiomycetes were predominant on sound berries, contrary to the observed predominance of *C. zemplinina* and *H. uvarum* when grape whole bunches were analysed. In particular, the apiculate *H. uvarum*/*K. apiculata* appears to be the most common grape berry species worldwide (Table 2) which is consistent with its predominance in the beginning of spontaneous must fermentations.

The proliferation of the third yeast group is explained by the high nutrient availability resulting from grape damage. Damaged grapes possess, besides much higher cell counts, wider species diversity than sound grapes (Table 3). Basidiomycetes may be still present in numbers similar to those of sound grapes but their proportion is strongly decreased by the proliferation of ascomycetous species (Barata et al., 2008a, 2008b, in press). *H. uvarum* and *C. zemplinina* may be present in higher numbers but their relative proportion also decreases in favour of the fermentative yeasts (e. g. *Pichia* spp., *Zygosaccharomyces* spp., *Zygoascus* spp., *Torulaspota* spp.), which may occasionally dominate the overall microbiota (Barata et al., 2008a, in press). Although regarded as oxidative or weakly fermentative, the physiological heterogeneity of *Candida* spp. and *Pichia* spp. explains the recovery of strongly fermentative species of these genera in damaged grapes or fermented juices (Barata et al., 2008b). In particular, with sour rot and honeydew, Barata et al. (2008a, 2008b, in press) proposed that *Zygoascus hellenicus*/*Candida steatolytica*, *P. terricola* and *P. kudriavzevii* (the two latter were named as *Issatchenkia* spp.) may be regarded as zymological indicators of these types of damage. *C. zemplinina* (formerly identified as *C. stellata*) is a typical contaminant of botrytised juice fermentations (Sipiczki, 2003) but its dissemination is also spread to sound grapes (Barata et al., 2008a, in press).

The proposed grouping is based on yeast species diversity. While acetic acid bacteria are clearly favoured by damage, it is not so for lactic acid bacteria and their grouping may be questionable. Grape lactic acid bacteria include *Lactobacillus* spp. and *Pediococcus* spp. mostly recovered after enrichment (Table 4). In particular, *O. oeni* has only

Table 4
Bacterial species detected in both sound and damaged grapes.

Bacterial groups	Species	References ^{a, b}	
Lactic acid	<i>Lactobacillus plantarum</i>	Bae et al. (2006); Lafon-Lafourcade et al. (1983); Renouf et al. (2005, 2007)	
	<i>L. hilgardii</i>	Lafon-Lafourcade et al. (1983)	
	<i>L. casei</i> , <i>L. sanfranciscensis</i>	Renouf et al. (2005, 2007)	
	<i>L. lindneri</i> , <i>L. kunkeei</i>	Bae et al. (2006)	
	<i>L. kefirii</i> ^c , <i>L. mali</i> ^c , <i>L. plantarum</i> ^c	Bae et al. (2006)	
	<i>L. brevis</i>	Prieto et al. (2007)	
	<i>Lactococcus lactis</i>	Bae et al. (2006); Francesca et al. (2010)	
	<i>Leuconostoc fallax</i>	Francesca et al. (2010)	
	<i>Lc. mesenteroides</i>	Renouf et al. (2005, 2007)	
	<i>Oenococcus oeni</i> ^a	Renouf et al. (2005, 2007)	
	<i>Pediococcus parvulus</i> , <i>P. damnosus</i> , <i>P. acidilactici</i>	Renouf et al. (2005, 2007)	
	<i>Weissella paramesenteroides</i>	Bae et al. (2006); Renouf et al. (2005, 2007)	
	Acetic acid	<i>Acetobacter aceti</i>	Barbe et al. (2001); González et al. (2005); Joyeux et al. (1984)
		<i>A. pasteurianus</i>	Barbe et al. (2001); Joyeux et al. (1984)
		<i>A. cerevisiae</i>	Prieto et al. (2007)
		<i>A. orleanensis</i> ^c , <i>A. syzygii</i> ^c	Barata et al. (in press)
		<i>Gluconobacter oxydans</i>	Barata et al. (in press); Barbe et al. (2001); González et al. (2005); Joyeux et al. (1984); Prieto et al. (2007); Renouf et al. (2005, 2007) Barata et al. (in press)
<i>G. cerinus</i>		Nisiotou et al. (2011)	
<i>Gluconoacetobacter hansenii</i>		Barata et al. (in press); González et al. (2005)	
<i>Gl. saccharivorans</i> , <i>Gl. intermedius</i> ^c		Barata et al. (in press)	
Other species		<i>Burkholderia vietnamiensis</i> , <i>Pseudomonas jessenii</i> , <i>Serratia rubidae</i> ,	Renouf et al. (2005, 2007)
		<i>Enterobacter gergoviae</i> , <i>Leifsonia xyli</i> , <i>Enterococcus faecium</i> ,	
	<i>Bacillus mycoides</i>		
	<i>Enterococcus durans</i> , <i>E. faecium</i> , <i>E. avium</i>	Bae et al. (2006)	
	<i>E. hermanniensis</i> ^c	Bae et al. (2006)	
	<i>Acinetobacter</i> spp., <i>Curtobacterium</i> spp., <i>Enterobacter</i> spp.,	Prieto et al. (2007)	
	<i>Stenotrophomonas maltophilia</i> <i>Serratia</i> spp., <i>Staphylococcus</i> spp.		
	<i>Enterococcus durans</i> , <i>Kocuria kristinae</i> , <i>Staphylococcus</i>		
	<i>saprophyticus</i>	Barata et al. (in press); Nisiotou et al. (2011)	
	<i>Klebsiella oxytoca</i> , <i>Bacillus subtilis</i> ^c , <i>Erwinia</i> spp., <i>Pantoea dispersa</i> ,	Nisiotou et al. (2011)	
	<i>Enterobacter ludwigii</i> ^c , <i>Tatumella ptyseos</i> , <i>Providencia rettgeri</i> ^c ,		
	<i>Serratia marcescens</i> ^c , <i>Citrobacter freundii</i> ^c		

^a Microbial detection after culture enrichment (Bae et al., 2006; Prieto et al., 2007; Renouf et al., 2005, 2007).

^b Microbial detection after plating (Barbe et al., 2001; Francesca et al., 2010; González et al., 2005; Joyeux et al., 1984; Lafon-Lafourcade et al., 1983; Renouf et al., 2005, 2007).

^c Recovery from damaged grapes only.

been recovered from grapes using methodologies adequate to elicit minority populations (Renouf et al., 2005, 2007). Then, the justification to include lactic acid bacteria in the first group is related with their rare isolation either from sound or damaged berry surfaces. In fact, these bacteria are regarded as anaerobic (or semi-anaerobic) and nutritively fastidious, probably unable to compete with yeasts and acetic acid bacteria under the aerobic conditions of the grape berry.

Acetic acid bacteria, dominated by *Gluconobacter* spp., but also *Acetobacter* spp., have been recovered from sound grapes (Table 4), and are particularly frequent in grey rotten grapes (Barbe et al., 2001). Barata et al. (in press) found that *Acetobacter* spp. are the dominant acetic acid bacteria in the final stage of sour rot, while *Gluconacetobacter* spp. and *Gluconobacter* spp. were mostly isolated from sound berries.

3.3. The case of *Saccharomyces* spp. and wine spoilage yeasts

The *Saccharomyces* spp. and wine spoilage species are mostly present in low numbers and in low frequencies, even in damaged grapes. Davenport (1973, 1974), studying English vineyards in 1965, carried out a thorough ecological study where the ecosystem was divided in atmosphere, phyllosphere and rhizosphere. *Schizosaccharomyces* spp. was found in rhizosphere (soil and leaf litter). *Z. baillii* was found in mummified fruit (grapes) in soil. *Z. rouxii* was found in mummified pears in soil from an adjacent orchard and from wild flowers. This author, by direct techniques, only found *S. cerevisiae* in one sample of acid soil, mushroom compost and vine flowers but not from mature fruit. The isolation of this species from grapes was only achieved by enrichment cultures. Subsequent works have confirmed this assumption (De la Torre et al., 1999; Rementeria et al., 2003; Renouf et al., 2005). Moreover, the frequency of occurrence of *S. cerevisiae* was found about 0.05 to 0.1% in sound berries and 25% in damaged berries, usually with numbers of about 10^5 – 10^6 /berry (Mortimer and Polsinelli, 1999). Interestingly, Schuller et al. (personal communication) recovered *S. cerevisiae* from auto-enrichment cultures using whole bunches with a frequency higher in Portuguese regions more prone to grape rot. The fact that *S. cerevisiae* was absent from all ferments of a hot and dry region (Alentejo) corroborates the importance of damage on its proliferation.

Wine spoilage species should behave similarly to *S. cerevisiae*, given their physiological similarities. Jolly et al. (2003), analysing bunches, reported *Z. baillii* as the predominant strain (>50%) in one site over 12, but no mention to damage was made. The genera *Zygosaccharomyces* spp. and *Torulospira* spp. were detected at higher frequencies in grapes affected by noble rot, sour rot and honeydew, suggesting their adaptation to conditions of reduced water activity and presence of weak organic acids (Barata et al., 2008a, 2008b; Nisioutou and Nychas, 2007). However, even in sour rot berries, they are rarely the dominant population probably because the aerobic conditions favour the fast growing populations of *Candida* spp., *Hanseniaspora* spp. and *Pichia* spp..

Concerning other spoilage species, their isolation from grapes is very rare. *Dekkera* spp. has been recovered from both sound and sour rot berries, apparently with higher numbers in rotten berries (Guerzoni and Marchetti, 1987), and *D. bruxellensis* from grapes after enrichment (Renouf and Lonvaud-Funel, 2007). In addition to the reports of Davenport (1973, 1974), *Sc. pombe* has been mentioned by Florenzano et al. (1977) and *Saccharomyces ludwigii* by Combina et al. (2005), but no relation to grape health was established. Future works with appropriate sampling plans are required to definitely assess their dissemination in the vineyard environment and track the sources of wine spoilage species.

3.4. Habitat shared by other microbial species

The microbiota of grapes also includes fungi (see Table 1) which may dominate under favourable weather conditions accompanied

by inefficient phytochemical utilisation. Fungal obligate parasites are able to penetrate through the intact grape skin by their own biochemical and mechanical activities and are responsible for high economic losses. The main species are the oomycete *Plasmopara viticola*, responsible for downy mildew, and the ascomycetes *Erysiphe necator* (powdery mildew), *Elsinoë ampelina* (anthracnose), *Guignardia bidwellii* (black rot) and *Pseudopezicula tracheiphila* (rotbrenner). Their biology and epidemiology, well-known by phytopathologists, are strongly dependent on weather conditions (mainly temperature and humidity). Berry susceptibility to these diseases decreases from berry set until véraison, after which development of ontogenic resistance explains the absence of parasite attacks, even in the absence of phytochemical treatments (Kennelly et al., 2005). Therefore, these populations are absent when the components of the WMC dominate the berry microbiota, but the induced damage may stimulate their growth as shown by the effect of diffuse powdery mildew (Kennelly et al., 2005).

Saprophytic moulds, inducing grape rots, include *B. cinerea* (grey rot), and other ubiquitous genera (e. g. *Cladosporium* spp., *Aspergillus* and *Penicillium* spp.) (Serra and Peterson, 2007), are especially visible when berries are injured, either within tight bunches or on bunch surfaces, competing with the WMC microbiota. The spores of these moulds are spread all over vine tissues and germinate when temperature and humidity are appropriate. Studies are required to enlighten possible interactions among overall grape microbiota and explain their succession and predominance during grape ripening. Probably, the emergence of sour rot against grey rot, besides better phytochemical efficiency against the latter, is related with higher average temperatures during ripening and harvest.

Ubiquitous bacterial species have also been reported on grapes as a result of contamination from the surrounding environment or, hypothetically, as constituents of endophytic populations (Table 4). The genus *Burkholderia* spp. (Renouf et al., 2005) is a representative of the latter while *Bacillus thuringiensis*, used as a biopesticide, may be predominant at harvest (Fleet, 2003). A recent report, revealed the presence of *Enterobacter ludwigii* after laboratorial must fermentation justifying further studies to understand its possible effect on wine quality (Nisioutou et al., 2011). On the contrary, Barata et al. (in press) did not find these environmental bacterial species on winery fermenting musts or wines.

4. Factors influencing species diversity

The microbial communities on grapes may be affected by a large number of factors as described for other fields of microbial ecology (Boddy and Wimpenny, 1992). In vineyards and winemaking, Pretorius et al. (1999) listed a series of variables that may affect the ecology of grape yeasts. Many of those variables are not independent and may be gathered in broad groups of effects, as described below. Moreover, all factors commonly described as influencing grape microbiota (rainfall, wind, temperature, diseases, pests, viticultural practises, etc.), affect primarily skin integrity and so their impact will be discussed further taking into account the expected changes induced by berry damage.

4.1. Climatic conditions

The climatic and microclimatic conditions include the effect of temperature, UV exposure, rainfall, sunlight and winds. Several studies mention that diversity and quantity of microbial populations are dependent on these conditions. However, results are often unclear because it is not easy to apply the scientific method. For instance, rainy vintages lead to higher use of phytochemicals, higher fungal proliferation and higher berry damage, in conjunction with lower UV irradiation. Concerning total yeast counts, Combina et al. (2005) found that years with increased rainfall yielded higher counts,

probably due to increase in berry volume allowing release of juice in joint areas such as the area between the pedicel and the berry, and higher exosmosis leading to nutrient release on the grape surface. With careful sound berry sampling, Čadež et al. (2010) also found that colder harvests with higher rainfall lead to higher yeasts counts. On the contrary, Comitini and Ciani (2006) found 10 times less total counts in years with high rainfall. Rementeria et al. (2003) also found higher numbers in warmer and drier years.

The reports on species diversity are also not conclusive. Some works state that oxidative yeast, like *M. pulcherrima* increase in years with increased rainfall (Combina et al., 2005; Longo et al., 1991; Poulard et al., 1981). Yanagida et al. (1992) found higher proportions of basidiomycetous yeasts in late crops from chilly climatic conditions while *K. apiculata* was more frequent in mild climates. Rementeria et al. (2003) isolated mostly *K. apiculata* in the warmer years but De La Torre (1999), in a hot region, never detected *K. apiculata* on grapes. Ganga and Martínez (2004) detected different proportions of non-*Saccharomyces* in two different vintages and in two producing regions. One harvest had higher rainfall and greater species variability was reported, but only in one region. The other showed less diversity that was explained by the use of fungicides against *B. cinerea*. Without noticeable fungicide effect, Čadež et al. (2010) found dominance by *A. pullulans* and lower species diversity in the warmer and drier vintage.

Large scale works do not demonstrate any relation between climatic conditions and yeast diversity. Jolly et al. (2003), in South Africa, during 3 vintages and in 4 different regions, did not find any pattern linking non-*Saccharomyces* species to climatic zones. Schuller et al. (2000) and Valero et al. (2005), in Northwest Portugal, from 2001 to 2003, using sensitive molecular and statistical techniques, found no correlation between *S. cerevisiae* and regional climatic conditions. Van der Westhuizen et al. (2000b) (1995 to 1998) and Khan et al. (2000) found different populations of *S. cerevisiae* in coastal and inner warmer regions of South Africa. These authors stated that, when a large number of vintages are analysed, appearance/reappearance cycles of certain strains have no obvious explanation.

4.2. Vineyard treatments

The main vineyard treatment studied is related with the use of pesticide treatments, mainly those against fungi (downy mildew, powdery mildew and grey rot). The studies are either based on analysing grapes after vineyard treatment, which do not exclude the influence of other factors, or from auto-enrichment fermentations which cannot be correctly extrapolated to evaluate the variations on berry microbiota.

Viviani-Nauer et al. (1995) found that pesticides decreased yeast population and diversity in fermenting musts. Cabras et al. (1999) reported the absence of effect on fermentation of *S. cerevisiae* by 6 different fungicides while fermentation by *K. apiculata* was stimulated. Guerra et al. (1999) concluded that pesticide affect diversity and frequency of *S. cerevisiae* and other species by comparing two different groups of pesticide application and one abandoned vineyard. However, careful analyses of results do not allow this conclusion. For instance, i) the abandoned vineyard had a diversity similar to one of the treated groups; ii) biodiversity was determined after fermentation and so any effect if existent should be on biodiversity of species enduring fermentation and not on grape contamination yeasts; and iii) in two other vintages, distribution of *Saccharomyces* was different for one grapevine variety and not for the other, perhaps due to its more resistant skin.

Ganga and Martínez (2004) detected less diversity of non-*Saccharomyces* species, which was explained by the use of fungicides against *B. cinerea*. However, while Ganga and Martínez (2004) did not find reduced *S. cerevisiae* numbers after fungicide use, Regueiro et al. (1993) and Valero et al. (2007) recovered lower numbers of this

species. Van der Westhuizen et al. (2000b), from 1995 to 1998, found a lower incidence of *S. cerevisiae* in 1997 where fungal infestation and chemical use were higher due to high rainfall, but clearly stated that a direct relation could not be made.

Longo et al. (1991) found that the number of oxidative yeasts increased in years with high rainfall, with high fungal proliferation, higher fungicide treatments and higher berry damage. Combina et al. (2005) found that oxidative yeasts were higher in years with increased rainfall (*M. pulcherrima* and *C. stellata* higher than *K. apiculata*) which were not attributed to chemical treatments because they were identical in both years. Čadež et al. (2010), with careful berry selection, showed that, after the safety interval, fungicides against *Botrytis* had a minor impact on the composition of grape berry microbiota and untreated grapes were less contaminated. Further, Oliva et al. (2007) did not find any adverse effect on yeast numbers and must fermentation, of several fungicides even when applied in the day of harvest.

When studying phytochemicals, attention should be paid to the product adjuvants as well. Rogiers et al. (2005) found that the spray adjuvants used to increase the effect of pesticides affect the epicuticular wax and favour *B. cinerea* infection and lower the contamination yeasts on grape surface. On the contrary, Čadež et al. (2010) attributed higher yeast counts on iprodione (an anti-*Botrytis* product) treated grapes to the presence of such adjuvants.

More recent works concern the differences from organic and conventional farming systems and it is tempting to conclude that organic farming leads to higher biodiversity, both in *S. cerevisiae* and in non-*Saccharomyces* yeasts (Cordero-Bueso et al., 2011). However, these authors recovered yeasts after blending 2 kg of grapes and using the juice as auto-enrichment medium that makes the use of the reported ecological indexes questionable. Moreover, their results may be explained by the effect of grape damage. In fact, *S. cerevisiae* was not detected in the ferments of two out of the 3 grape varieties in the 3 analysed vintages when pesticides were used, while it was always found in all samples from organic farming. Additionally, some of the recovered species from both farming systems (*C. sorbosa/H. occidentalis*, *C. stellata*, *K. thermotolerans*, *T. delbrueckii*) are consistent with those recovered from damaged grapes (see Table 2). Then, if winemakers want to obtain grapes with biodiversity similar to that of organic farming, the solution might be to leave some grapes rotting and use it in the fermentations.

4.3. Biotic factors

4.3.1. Microbial vectors

The association between yeasts and invertebrates is well-known in general ecological studies (review of Ganter, 2006). Miller and Phaff (1962) showed that figs, infected by wasps and *Drosophila* (in external body parts), carry apiculate yeasts and *C. stellata*. Few data available from vineyards showed that bees and wasps carry yeasts for the grapes (Davenport, 1973, 1974; Stevic, 1962). Fermaud et al. (2000), studying sour rot, associated *Drosophila* flies with *K. apiculata*, *C. stellata*, *P. membranifaciens*, *M. pulcherrima*, *C. krusei* and acetic acid bacteria. Moreover, yeasts and acetic acid bacteria, carried by *Drosophila* flies were found essential to trigger sour rot (Barata et al., accepted for publication). Lactic acid bacteria, mostly *L. plantarum*, were isolated from the intestinal tract of *Drosophila simulans* collected from a winery (Groenewald et al., 2006).

Regarding birds, a recent survey evidenced that swabs of bird beaks and the initial part of the digestive tract are reservoirs of several grape contamination species, like *H. uvarum* (Francesca et al., 2010).

4.3.2. Microbial interactions and enzymatic activities

There are few reports on the interactions between microbial populations that possibly include killer toxin, antibiotic and quorum sensing mechanisms (Fleet, 2003; Golubev, 2006). The interactions

between resident populations may affect diversity but there is no clear explanation. Yeast–yeast interactions are mostly studied regarding the killer effect, but this activity is probably not relevant in natural populations (Sangorrín et al., 2001). In particular, Van der Westhuizen et al. (2000b) found that killer activity did not affect the distribution of *S. cerevisiae*. The yeast like fungi *A. pullulans* seems to reduce basidiomycete diversity but results require confirmation (Čadež et al., 2010). Research directed to table grape storage showing inhibition of moulds by particular yeast species may provide some clues to future works on natural environments (Golubev, 2006).

Enzymatic activities may favour the survival on grape surface but the real effect remains to be elucidated. Moore et al. (1988) found lipolytic activity in all strains studied, suggesting a competitive advantage on the cutin of grape surfaces. Khan et al. (2000) also observed that all grape isolates exhibited identical results in the hydrolysis of various compounds: cellulose negative, starch negative, poligalacturonase negative, cellobiose positive, β -glucan negative and arbutin positive. Perhaps, these traits are common to most grape contaminants and so do not constitute competitive advantages among yeast species. Gognies et al. (2006) showed that *S. cerevisiae* is a phytopathogen to grapevine plantlets *in vitro*. The mechanisms depend on the filamentous growth process and the endopolygalacturonase activity alone was not responsible for invasion of plant tissues. This ability to form pseudohyphae by *S. cerevisiae* may enable the penetration in tissues and allow persistence (Khan et al., 2000), but results have not been validated under field conditions.

4.4. Geographic location and vineyard factors: age, size, grape variety and vintage year

There is a broad line of thought considering that the grape microflora is dependent on the vineyard location, grape variety and other vineyard related factors. The wish for this belief is that wine tipicity may be, at least partially, dependent on the specific grape microbiota of the producing region. The ultimate goal is to find yeasts according to each wine “terroir”. We believe that the available data do not allow conclusions on the existence of “terroir” species/strains, as described below. Works concluding otherwise are probably based on insufficient sample numbers, not accounting for variability of microbial populations in space and time.

With sound grapes, Guerzoni and Marchetti (1987) could not explain the variability within the same year for different varieties and within the same variety for different years. Poulard et al. (1981) concluded that the grape variety did not influence microbiota composition. De La Torre et al. (1999) found that the diversity varied with the vintage year. Sabate et al. (2002) found in one grape variety mostly basidiomycetous (similar to that of soil, bark and leaves) and, in the other, *H. uvarum* and *C. zeylanoides* predominated, but the same grape variety in the same vineyard showed a different species proportion in two different vintages. Zahavi et al. (2002) found lower numbers in one cultivar but the diversity was not dependent on the cultivar. Rementeria et al. (2003) found no correlation between grape variety or origin of grapes and number of yeasts. Chamberlain et al. (1997), in icewine grapes, found a lack of yeast species reproducibility in samples taken at different times or different parts of the same vineyard.

The most thorough studies of yeast distribution in vineyards concern *S. cerevisiae* and the conclusion may be conceivably extended to wine spoilage species. Van der Westhuizen et al. (2000a), Khan et al. (2000), Schuller et al. (2005), Schuller and Casal (2007) and Valero et al. (2005, 2007), on different hemispheres, found essentially the same results. They observed that indigenous *S. cerevisiae* populations on grapes were subjected to natural fluctuations of periodical appearance/disappearance, but no attempt to define “terroir” strains was made because no strain common to all sites, in one region, was found. Furthermore, the studies of Schuller et al. (2005), Schuller and Casal (2007) and Valero et al. (2005, 2007) also concluded that

S. cerevisiae from commercial starters were only detected near water running off the winery and that there was no influence of starter utilisation in the biodiversity of *S. cerevisiae* populations in the vineyards. Pretorius et al. (1999), describing close vineyards with equal climate, stated that intra-annual variations should be attributed to other vineyard factors like age and size. The vineyard age and size were not evaluated by Valero et al. (2007) but, from 2001 to 2003, found greater biodiversity in larger and older wine regions.

The saprophyte mould *B. cinerea* is known to overwinter in vine tissues (Gabler et al., 2003) and wine spoilage yeasts, also of saprophytic nature, could conceivably behave in the same manner. However, this hypothesis has not been demonstrated. Therefore, yeasts must be vectored from the surroundings to berry surface and be able to thrive afterwards, mainly during the period between *véraison* and harvest (about 2 months long) when the cuticle is thinner and VOCs attracting insects are produced. Colonisation should be driven by vector dissemination and further growth, upon inoculation, depends on the nutrient availability on berry skin. As a result, the grape berry is under the overwhelming influence of the surrounding environment from which vectors bring all kinds of yeast species. If the berry is sound, the usual plant colonisers, basidiomycetes and *A. pullulans*, predominate. If berries are damaged, the influence of environment is overcome and the predominance is diverted to yeast species rarely found in nature, as described before. Under this view, it is difficult to accept that the microbiota of a certain grapevine may be restrained by regional boundaries coincident with a particular “terroir”.

5. Influence of grape damage or overripeness on grape juice and wine quality

The detrimental influence of damaged grapes on quality is long known by enologists, being widely accepted that when damaged grapes arrive at the winery the resulting wine will be of lower rate. Surprisingly, the mechanisms explaining the detrimental effect of grape damage are not well understood. Apparently grape juice composition is affected in a negative way giving rise to unbalanced wines, but the origin of molecules responsible for the disorder or the role of fermentative microbiota are not fully understood. However, available data highlight the main role of fungal pathogens or saprophytes on wine depreciation while yeasts and bacteria are mostly regarded as innocuous contaminants. The two most famous exceptions to this quality depreciation are the wines produced from dried grapes—e.g. *Passito* wines, *vins de paille*—and the highly valued late harvest wines, produced with grapes affected by the noble rot mould *Botrytis cinerea*—e.g. Tokay, Sauternes.

5.1. Dried or overripe grapes

Grapes are naturally or artificially dried to obtain certain types of wine (*Passito*, *vin de paille*) or become dried due to unwanted overmaturation. In this kind of grapes it is expectable to find osmotolerant species. Davenport (1974) found *Z. bailii* and *Z. rouxii* in overripe grapes. Caridi and Audino (1997) isolated few *Z. bailii* from dried grapes that were not present in immature or mature grapes, being *H. guilliermondii*, the most frequent species in all 3 ripening stages. In mature and dried grapes *S. cerevisiae* appeared, but strains (former races *aceti*, *capensis*, *globosus*) were considered as having not good enological properties. Dried grapes had more diversity, because *Candida* spp. also appeared (Caridi and Audino, 1997). In other studies, no particular differences between overripe and sound grapes were observed (Prakitchaiwattana et al., 2004). Knowing that this particular type of wine requires dried grapes it is expected that this grape microbiota has no harmful effect on quality. Possible concerns are related with the occurrence of stuck fermentations due to high initial sugar content. When fungal growth accompanies berry dehydration

changes may occur in aroma composition (Noble-rot) or in the concentration of mycotoxins (Malfeito-Ferreira et al., 2009).

5.2. Noble-rot grapes

Noble-rot or late-harvest wines owe their quality to particular climate conditions that enable the drying process to be accompanied by desirable *B. cinerea* activities. These grapes bear higher constant microbial numbers (yeast and bacteria) during 1 week before maturation than healthy grapes (Barbe et al., 2001). Magyar and Bene (2006) listed the species isolated from these grapes during 5 years, showing that *C. pulcherrima*, *H. uvarum* and basidiomycetous yeasts (*Sporidiobolus*, *Rhodotorula*, *Cryptococcus* and *Filobasidium* spp.) were the most frequent species. There was only one isolate from grapes belonging to *Brettanomyces nanus* or to *Zygosaccharomyces* spp. The species dominant in the winery was *C. zemplinina*, already detected in other studies (Mills et al., 2002; Sipiczki, 2003), but without clear technological significance. It seems to be able to decrease volatile acidity (Cocolin et al., 2011b) that is a favourable ability.

The volatile compounds identified in sweet botrytised wines (e. g. Tokay, Fiano, Sauternes) belong to several chemical classes (e.g. lactones, terpenes, aldehydes, ketones, furanones) (Genovese et al., 2007; Sarrazin et al., 2007 and references cited therein).

The mould affects grapes by a smooth drying process, but the major benefit seems to be the alteration of grape metabolism by *Botrytis* that stimulates the production of a S-glutathione conjugate which gives S-3-(hexan-1-ol)cysteine (P-3SH) in musts. This molecule is the cysteinylated precursor of the most abundant volatile thiol in wine (3-sulfanylhexanol, 3SH) (Thibon et al., 2009). Therefore, yeasts should play a minor role in the particular aromatic features of these wines, if any.

5.3. Grey rot

The most common and feared grape rot—grey rot—is due to *B. cinerea* growing on grapes favoured by high rainfall before or during vintage. The fungal strong oxidasic and esterase activities are responsible for deterioration of phenolic compounds (anthocyanins, hydroxycinnamic acids and flavanols), for transformation of terpenes into less odourous compounds and for hydrolysis of ethyl esters of fatty acids (La Guerche et al., 2006). Moreover, several studies have shown that grape rot, due to the association of *B. cinerea* with other, less visible, fungi (*Penicillium* spp., *Rhizopus* spp.) frequently leads to the development of organoleptic defects in grapes and wines. These compounds have been identified as 2-methylisoborneol, (–)-geosmin, 1-octen-3-one, 1-octen-3-ol, 2-octen-1-ol, and 2-heptanol (La Guerche et al., 2006 and references therein). This mould also induces the production of a pathogenesis-related (PR) protein causing haziness in white wines (Girbau et al., 2004).

Yeasts and bacteria have been isolated from grey rot grapes (see Table 2) but their role in decreasing wine quality should be irrelevant when compared with the detrimental effects of *B. cinerea*.

5.4. Powdery mildew

Powdery mildew (oidium) due to the fungus *Uncinula necator* is an important disease for the vineyard affecting grape yield, juice and wine quality (Calonnet et al., 2004), titratable acidity (TA), total phenolics, hydroxycinnamates and flavonoids (Stummer et al., 2005). It leads to the occurrence of a very characteristic and sometimes intense mushroom-type odour default (Darriet et al., 2002; Stummer et al., 2005). Strongly odourant compounds were identified as 1-octen-3-one (mushroom odour), (Z)-1,5-octadien-3-one (geranium-leaf odour). During alcoholic fermentation, the enzymatic reduction of 1-octen-3-one and (Z)-1,5-octadien-3-one to much less odourant compounds, namely 3-octanone and (Z)-5-octen-3-one,

was observed (Darriet et al., 2002). Those results explain to some extent the disappearance of the fungal aroma specific to powdery mildew grapes during alcoholic fermentation. The influence on wine odour is very effective being noticed in juice and wines with as little as 1–5% of rotten bunches (Stummer et al., 2005), but the partners of the wine microbial consortium have no noticeable effect. The oidium also induces the formation of PR proteins, as mentioned before concerning grey rot (Girbau et al., 2004).

5.5. Diffuse powdery mildew (DPM)

Diffuse powdery mildew (DPM) is the initial stage of powdery mildew when the damage is not visible to the naked eye. It induces strong changes in microbiota composition. The difference between healthy and affected grapes is between a total count of 10^2 – 10^4 to about 10^4 – 10^7 /ml, including yeast, lactic and acetic bacteria (Gadoury et al., 2007). These microorganisms increase volatiles (ethanol, ethyl acetate, acetic acid) that attract insects (sap beetles, ants, wasps, yellow jackets) which damage berry and promote bunch rot severity by creating entry points to spoilage organisms. It is conceivable that grey or sour rot may follow the initial infection by DPM, but it is difficult to evaluate the detrimental effect of this initial step of powdery mildew *per se*.

5.6. Sour rot

The description of sour rot disease is somewhat unclear perhaps because it occurs simultaneously with other diseases, particularly when mould growth is evident. Formerly, it was regarded as the final stage of grey rot given the observation of *Botrytis* in affected berries (Bisiach et al., 1986). Other authors related it to the activity of saprophytic fungi like *Aspergillus carbonarius* (Dimakopoulou et al., 2008) or *Colletotrichum acutatum* (Meunier and Steel, 2009), and to *Acetobacter* spp. (Oliva et al., 1999). The diseases named as bunch or ripe rot may be taken as synonyms, given the analogy to the sour rot symptoms (Loinger et al., 1977; Meunier and Steel, 2009). The massive presence of yeasts, in the absence of moulds or bacteria, is considered as a requirement for the development of the disease (Marchetti et al., 1984). Blancard et al. (2000) found that the most effective agents, inducing symptoms *in vitro* and *in vivo* were the yeasts *K. apiculata* and *C. stellata* and the acetic acid bacteria *Gluconobacter* spp. and *A. pasteurianus*. It seems that, without acetic acid bacteria, disease symptoms are minor (Barata et al., accepted for publication). Probably, the best way to describe the microbial effects of sour rot is as the result of a spontaneous fermentation of juice, released on berry surface, by a natural mixed culture of yeasts and acetic acid bacteria, where the major final product with impact in wine quality is acetic acid (Barata et al., accepted for publication).

The effect on winemaking has been scarcely studied, being responsible for yield losses compensated by higher initial sugar and colour concentration (Loinger et al., 1977). However, colour is not stable and wine shows higher levels of volatile acidity and residual sugars (Barata et al., 2011b).

Guerzoni and Marchetti (1982) regarded acetic acid as the chemical indicator of the disease. Marchetti et al. (1984) studied the volatiles produced by the yeasts and observed that species *M. pulcherrima* and *H. uvarum* were the single species yeasts providing volatiles (mainly ethanol and ethyl acetate) similar to those of rotten grapes and absent from sound or *B. cinerea* infected grapes. According to these authors *Endomycopsella* spp. (now *Saccharomyopsis* spp.) enabled tissue penetration, but did not produce volatile acidity or ethyl acetate, then *M. pulcherrima* and *H. uvarum*, produce it and spoil the grape. The wine spoilage genera *Brettanomyces* spp. and *Zygosaccharomyces* spp. were found in both grapes, but more frequently in sour rotten samples.

Recently, phenylacetic acid and ethyl phenylacetate were found as key aroma components of grapes affected by sour rot (Barata et al., 2011a). These authors speculated that phenylacetic acid is produced on berry skin tissue as a plant defence mechanism, being esterified to ethyl phenylacetate during fermentation and storage. Furthermore, they observed a loss in the fatty acids and their ethyl esters which was attributed to the consumption of precursor amino acids by the contaminant microflora on damaged grapes.

5.7. Other forms of damages

Wine quality may be affected by other forms of damage which have received little attention in the literature. Recently, the effect of *Cladosporium* rot was reported in delayed harvests in Chile (Briceño et al., 2009). This type of rot reduced colour, aroma, and flavour in Cabernet Sauvignon and Carménère wines but there was no mention of yeast and bacterial activity.

Colletotrichum acutatum and *Greeneria uvicola* have been associated with ripe rot and bitter rot, respectively, in sub-tropical regions (Steel et al., 2007), decreasing grape quality and imparting wine taints (Meunier and Steel, 2009). Their emergence, with a concurrent decrease of grey rot, is probably related with climate changes (Steel et al., 2011).

Grape infection with *Aspergillus carbonarius* can increase the sugar and the citric acid concentrations (Leong et al., 2006) possibly affecting the perceptions of astringency and bitterness (Gawel, 1998; Noble, 1998).

Another two cases, not originally associated to grape health or yeast activity, may be the result of grape damage. One is attributed to the untypical “sweet-like” off-odour of Italian Aglianico del Vulture wines, due to ethyl phenylacetate, associated to late-ripening and empirically connected to some cryptogamic diseases (Tat et al., 2007). The other, is the so-called “premature red wine ageing”, where the key compound was 3-methyl-2,4-nonanedione originated from a furan fatty acid (FFA) (Pons et al., 2008). This molecule is a widespread component of plant lipids (Sigrist et al., 2001), present in cutin (Velíšek and Cejpek, 2006) and related with plant defence mechanisms (Batna and Spiteller, 1994). It is conceivable that FFA and phenylacetic acid are produced in grape berry as a response to berry injury, as mentioned for sour rot. Increased levels of these aroma precursors in grape juices originate higher levels of aromatic compounds—3-methyl-2,4-nonanedione and ethyl phenylacetate—that affect wine flavour during storage.

Other concerns are related with microbial metabolites affecting human health. The agents are mostly saprophytic moulds of the genus *Aspergillus* producing ochratoxin A (as reviewed by Malfeito-Ferreira et al., 2009). Recently, fumonisin B2 produced by *A. niger* was reported as an additional potential risk in wine production (Logrieco et al., 2011). As far as we are aware wine sensory attributes are not affected by these mycotoxins.

6. Future prospects

The knowledge of microbial ecology in the vineyard environment has many gaps and it is possible to suggest several subjects of future research.

The first issue is concerned with the origin and dissemination of the main agents of wine fermentation—*S. cerevisiae* and *O. oeni*—and of the other components of the WMC. Despite all information available, it is not yet possible to state where these microorganisms come from and their persistence in the environment during all year is not clarified. Lachance (2003) speculated that wild *S. cerevisiae* are associated with an interface involving *Drosophila* spp., oaks and the surrounding soil. Probably, wine spoilage and bacterial species share the same sites. The microbial vectors (e. g. insects, birds, rain, dusts) are also poorly studied. Which is their relative importance in

berry colonisation? Which factors determine their attraction to the sources of microorganisms and to grape berries? Where do microorganisms overwinter? All these questions can only be solved when the overall environment gathering vine plants, microorganisms and vectors will be studied by multidisciplinary research teams, including vine physiologists, entomologists and microbiologists.

Acknowledgements

A. Barata was a recipient of a fellowship support by the Portuguese research foundation FCT (grant SFRH/BD/28451/2006). We are grateful to Dr. Sally Burr for revising the English language.

References

- Andorrà, I., Landi, S., Mas, A., Guillaumon, J.M., Esteve-Zarzoso, B., 2008. Effect of oenological practices on microbial populations using culture-independent techniques. *Food Microbiology* 25, 849–856.
- Arena, M.P., Romano, A., Capozzi, V., Beneduce, L., Ghariani, M., Grieco, F., Lucas, P., Spano, G., 2011. Expression of *Lactobacillus brevis* IOEB 9809 tyrosine decarboxylase and agmatine deiminase genes in wine correlates with substrate availability. *Letters in Applied Microbiology* 53, 395–402.
- Arroyo-López, F.N., Salvadó, Z., Tronchoni, J., Guillaumon, J.M., Barrio, E., Querol, A., 2010. Susceptibility and resistance to ethanol in *Saccharomyces* strains isolated from wild and fermentative environments. *Yeast* 27, 1005–1015.
- Bae, S., Fleet, G.H., Heard, G.M., 2006. Lactic acid bacteria associated with wine grapes from several Australian vineyards. *Journal of Applied Microbiology* 100, 712–727.
- Baffi, M., dos Santos Bezerra, C., Arévalo-Villena, M., Isabel Briones-Pérez, A., Gomes, E., Da Silva, R., 2011. Isolation and molecular identification of wine yeasts from a Brazilian vineyard. *Annals of Microbiology* 61, 75–78.
- Barata, A., González, S., Malfeito-Ferreira, M., Querol, A., Loureiro, V., 2008a. Sour rot-damaged grapes are sources of wine spoilage yeasts. *Fems Yeast Research* 8, 1008–1017.
- Barata, A., Seborro, F., Belloch, C., Malfeito-Ferreira, M., Loureiro, V., 2008b. Ascomycetous yeast species recovered from grapes damaged by honeydew and sour rot. *Journal of Applied Microbiology* 104, 1182–1191.
- Barata, A., Campo, E., Malfeito-Ferreira, M., Loureiro, V.I., Cacho, J., Ferreira, V., 2011a. Analytical and sensorial characterization of the aroma of wines produced with sour rotten grapes using GC–O and GC–MS: identification of key aroma compounds. *Journal of Agricultural and Food Chemistry* 59, 2543–2553.
- Barata, A., Pais, A., Malfeito-Ferreira, M., Loureiro, V., 2011b. Influence of sour rotten grapes on the chemical composition and quality of grape must and wine. *European Food Research and Technology*.
- Barata, A., Santos, S., Malfeito-Ferreira, M., Loureiro, V., accepted for publication. The role of the insect *Drosophila* sp. and of the grape berry microbiota in the development of sour rot disease. *Microbial Ecology*.
- Barata, A., Malfeito-Ferreira, M., Loureiro, V., in press. Changes in sour rotten grape berry microbiota during ripening and wine fermentation. *International Journal of Food Microbiology*.
- Barbe, J.C., de Revel, G., Joyeux, A., Bertrand, A., Lonvaud-Funel, A., 2001. Role of botrytized grape micro-organisms in SO₂ binding phenomena. *Journal of Applied Microbiology* 90, 34–42.
- Barnett, J.A., Winch, B., Delaney, M.A., Magson, A.B., Jones, E., 1972. Numbers of yeasts associated with wine grapes of Bordeaux. *Archiv für Mikrobiologie Arch Mikrobiol* 83 (1), 52–55.
- Barnett, J.A., Payne, R.W., Yarrow, D., 1990. *Yeast: Characteristics and Identification*, 2nd ed. Cambridge University Press, London, UK.
- Batna, A., Spiteller, G., 1994. Oxidation of furan fatty acids by soybean lipoxygenase-1 in the presence of linoleic acid. *Chemistry and Physics of Lipids* 70, 179–185.
- Belin, J.M., 1972. Recherches sur la répartition des levures à la surface de la grappe de raisin. *Vitis* 11 (2), 135–145.
- Bisiach, M., Minervini, G., Zerbetto, F., 1986. Possible integrated control of grapevine sour rot. *Vitis* 25, 118–128.
- Blancard, D., Grivot, E., Jailloux, F., Fermaud, M., 2000. Etiology of sour rot in vineyards located in South-West of France. IOBC/wprs Bulletin “Integrated Control in Viticulture”, 23, pp. 51–54.
- Boddy, L., Wimpenny, J.W.T., 1992. *Ecological Concepts in Food Microbiology*. *Journal of Applied Bacteriology Symposium Supplement*.
- Boundy-Mills, K., 2006. Methods for investigating yeast biodiversity. In: Péter, G., Rosa, C. (Eds.), *Biodiversity and Ecophysiology of Yeasts*. Springer, Berlin Heidelberg, pp. 67–100.
- Briceño, E.X., Latorre, B.A., Bordeu, E., 2009. Effect of *Cladosporium* rot on the composition and aromatic compounds of red wine. Efecto de la cladosporiosis en la composición y compuestos aromáticos de vinos tintos 7, 119–128.
- Cabras, P., Angioni, A., Garau, V.L., Pirisi, F.M., Farris, G.A., Madau, G., Emonti, G., 1999. Pesticides in fermentative processes of wine. *Journal of Agricultural and Food Chemistry* 47, 3854–3857.
- Čadež, N., Zupan, J., Raspor, P., 2010. The effect of fungicides on yeast communities associated with grape berries. *Fems Yeast Research* 10, 619–630.
- Calonne, A., Cartolar, P., Poupot, C., Dubourdieu, D., Darriet, P., 2004. Effects of *Uncinula necator* on the yield and quality of grapes (*Vitis vinifera*) and wine. *Plant Pathology* 53, 434–445.

- Capozzi, V., Ladero, V., Beneduce, L., Fernández, M., Alvarez, M.A., Benoit, B., Laurent, B., Grieco, F., Spano, G., 2011. Isolation and characterization of tyramine-producing *Enterococcus faecium* strains from red wine. *Food Microbiology* 28, 434–439.
- Caridi, A., Audino, P., 1997. Evoluzione della microflora lievitifforme durante la maturazione e la passitura delle uve. *Rivista di Viticoltura e di Enologia* 50, 3–8.
- Chamberlain, G., Husnik, J., Subden, R.E., 1997. Freeze-desiccation survival in wild yeasts in the bloom of icewine grapes. *Food Research International* 30, 435–439.
- Chavan, P., Mane, S., Kulkarni, G., Shaikh, S., Ghormade, V., Nerkar, D.P., Shouche, Y., Deshpande, M.V., 2009. Natural yeast flora of different varieties of grapes used for winemaking in India. *Food Microbiology* 26, 801–808.
- Ciani, M., Faticenti, F., Mannazu, I., 2002. Yeasts in winemaking biotechnology. In: Ciani, M. (Ed.), *Biodiversity and Biotechnology of Wine Yeasts*. Research Signpost, Kerala, India, pp. 112–123.
- Clavijo, A., Calderón, I.L., Paneque, P., 2010. Diversity of *Saccharomyces* and non-*Saccharomyces* yeasts in three red grape varieties cultured in the Serranía de Ronda (Spain) vine-growing region. *International Journal of Food Microbiology* 143, 241–245.
- Cocolin, L., Bisson, L.F., Mills, D.A., 2000. Direct profiling of the yeast dynamics in wine fermentations. *FEMS Microbiology Letters* 189, 81–87.
- Cocolin, L., Campolongo, S., Alessandria, V., Dolci, P., Rantsiou, K., 2011a. Culture independent analyses and wine fermentation: an overview of achievements 10 years after first application. *Annals of Microbiology* 61, 17–23.
- Cocolin, L., Dolci, P., Rolle, L., Rantsiou, K., 2011b. *Candida zemplinina*: approcci biotecnologici innovativi per la riduzione del contenuto in acido acetico in fermentazioni di mosti ad alto contenuto zuccherino. Workshop Qualità, Innovazione e Territorio: La Ricerca Microbiologica per i Vini del Futuro. P.R.C. Repubbliche srl, Palermo, Castello Utveggi, pp. 13–16.
- Combina, M., Mercado, L., Borgo, P., Elia, A., Jofré, V., Ganga, A., Martínez, C., Catania, C., 2005. Yeasts associated to Malbec grape berries from Mendoza, Argentina. *Journal of Applied Microbiology* 98, 1055–1061.
- Comitini, F., Ciani, M., 2006. Survival of inoculated *Saccharomyces cerevisiae* strain on wine grapes during two vintages. *Letters in Applied Microbiology* 42, 248–253.
- Cordero-Bueso, G., Arroyo, T., Serrano, A., Tello, J., Aporta, I., Vélez, M.D., Valero, E., 2011. Influence of the farming system and vine variety on yeast communities associated with grape berries. *International Journal of Food Microbiology* 145, 132–139.
- Darriet, P., Pons, M., Henry, R., Dumont, O., Findeling, V., Cartolaro, P., Calonne, A., Dubourdieu, D., 2002. Impact odorants contributing to the fungus type aroma from grape berries contaminated by powdery mildew (*Uncinula necator*); incidence of enzymatic activities of the yeast *Saccharomyces cerevisiae*. *Journal of Agricultural and Food Chemistry* 50, 3277–3282.
- Davenport, R.R., 1973. Vineyard yeasts—an environmental study. In: Board, R.G., Lovelock, D.H. (Eds.), *Sampling—Microbiological Monitoring of Environments*. Academic Press, London, pp. 143–174.
- Davenport, R.R., 1974. Microecology of yeasts and yeast like organisms associated with an English vineyard. *Vitis* 13, 123–130.
- De La Torre, M., Millan, M., Perez-Juan, P., Morales, J., Ortega, J., 1999. Indigenous yeasts associated with two *Vitis vinifera* grape varieties cultured in southern Spain. *Microbios* 100, 27–40.
- de Llanos, R., Fernández-Espinar, M.T., Querol, A., 2004. Identification of species of the genus *Candida* by analysis of the 5.8S rRNA gene and the two ribosomal internal transcribed spacers. *Antonie van Leeuwenhoek* 85, 175–185.
- De Vero, L., Gala, E., Gullo, M., Solieri, L., Landi, S., Giudici, P., 2006. Application of denaturing gradient gel electrophoresis (DGGE) analysis to evaluate acetic acid bacteria in traditional balsamic vinegar. *Food Microbiology* 23, 809–813.
- Deák, T., Beuchat, L.R., 1996. *Handbook of Food Spoilage Yeasts* CRC Press, Boca Raton, Florida.
- Di Maro, E., Ercolini, D., Coppola, S., 2007. Yeast dynamics during spontaneous wine fermentation of the Catalanesca grape. *International Journal of Food Microbiology* 117, 201–210.
- Dimakopoulou, M., Tjamos, S.E., Antoniou, P.P., Pietri, A., Battilani, P., Avramidis, N., Markakis, E.A., Tjamos, E.C., 2008. Phyllosphere grapevine yeast *Aureobasidium pullulans* reduces *Aspergillus carbonarius* (sour rot) incidence in wine-producing vineyards in Greece. *Biological Control* 46, 158–165.
- Esteve-Zarzoso, B., Belloch, C., Uruburu, F., Querol, A., 1999. Identification of yeasts by RFLP analysis of the 5.8S rRNA gene and the two ribosomal internal transcribed spacers. *International Journal of Systematic Bacteriology* 49, 329–337.
- Farris, G.A., Zara, S., Pinna, G., Budroni, M., 2002. Genetic aspect of flor yeasts Sardinian strains, a case of study. In: Ciani, M. (Ed.), *Biodiversity and Biotechnology of Wine Yeasts*. Research Signpost, Kerala, India, pp. 71–83.
- Fermand, M., Gravot, E., Blancard, D., Jailloux, F., Stockel, J., 2000. Association of *Drosophila* with microorganisms in Bordeaux vineyards affected by sour rot. IOBC/wprs Bulletin “Integrated Control in Viticulture”, 23, pp. 55–58.
- Fernández-Espinar, M., Esteve-Zarzoso, B., Querol, A., Barrio, E., 2000. RFLP analysis of the ribosomal internal transcribed spacers and the 5.8S rRNA gene region of the genus *Saccharomyces*: a fast method for species identification and the differentiation of flour yeasts. *Antonie van Leeuwenhoek* 78, 87–97.
- Fernández-Espinar, M.T., Llopis, S., Querol, A., Barrio, E., 2011. Molecular identification and characterization of wine yeasts. In: Carrascosa, A.V., Muñoz, R., González, R. (Eds.), *Molecular Wine Microbiology*, 1st ed. Elsevier, Academic Press, London, UK, pp. 111–141.
- Fleet, G.H., 2003. Yeast interactions and wine flavour. *International Journal of Food Microbiology* 86, 11–22.
- Fleet, G., Prakitichaiwattana, C., Beh, A., Heard, G., 2002. The yeast ecology of wine grapes. In: Ciani, M. (Ed.), *Biodiversity and Biotechnology of Wine Yeasts*. Research Signpost, Kerala, India, pp. 1–17.
- Florenzano, G., Balloni, W., Materassi, R., 1977. Contributo alla ecologia dei lieviti *Schizosaccharomyces* sulle uve. *Vitis* 16, 38–44.
- Fonseca, Á., Inácio, J., 2006. Phylloplane yeasts. In: Péter, G., Rosa, C. (Eds.), *Biodiversity and Ecophysiology of Yeasts*. Springer, Berlin Heidelberg, pp. 263–301.
- Francesca, N., Chiurazzi, M., Romano, R., Aponte, M., Settanni, L., Moschetti, G., 2010. Indigenous yeast communities in the environment of “Rovello bianco” grape variety and their use in commercial white wine fermentation. *World Journal of Microbiology and Biotechnology* 26, 337–351.
- Francesca, N., Settanni, L., Sannino, C., Aponte, M., Moschetti, G., 2011. Ecology and technological capability of lactic acid bacteria isolated during Grillo grape vinification in the Marsala production area. *Annals of Microbiology* 61, 79–84.
- Gabler, F.M., Smilanick, J.L., Mansour, M., Ramming, D.W., Mackey, B.E., 2003. Correlations of morphological, anatomical, and chemical features of grape berries with resistance to *Botrytis cinerea*. *Phytopathology* 93, 1263–1273.
- Gadoury, D.M., Seem, R.C., Wilcox, W.F., Henick-Kling, T., Conterno, L., Day, A., Ficke, A., 2007. Effects of diffuse colonization of grape berries by *Uncinula necator* on bunch rots, berry microflora, and juice and wine quality. *Phytopathology* 97, 1356–1365.
- Ganga, M.A., Martínez, C., 2004. Effect of wine yeast monoculture practice on the biodiversity of non-*Saccharomyces* yeasts. *Journal of Applied Microbiology* 96, 76–83.
- Ganter, P.F., 2006. Yeast and invertebrate associations. In: Péter, G., Rosa, C. (Eds.), *Biodiversity and Ecophysiology of Yeasts*. Springer, Berlin Heidelberg, pp. 303–370.
- Gawel, R., 1998. Red wine astringency: a review. *Australian Journal of Grape and Wine Research* 4, 74–95.
- Genovesse, A., Gambuti, A., Piombino, P., Moio, L., 2007. Sensory properties and aroma compounds of sweet Fiano wine. *Food Chemistry* 103, 1228–1236.
- Girbau, T., Stummer, B.E., Pocock, K.F., Baldock, G.A., Scott, E.S., Waters, E.J., 2004. The effect of *Uncinula necator* (powdery mildew) and *Botrytis cinerea* infection of grapes on the levels of haze-forming pathogenesis-related proteins in grape juice and wine. *Australian Journal of Grape and Wine Research* 10, 125–133.
- Giudici, P., Pulvirenti, A., 2002. Molecular methods for identification of wine yeasts. In: Ciani, M. (Ed.), *Biodiversity and Biotechnology of Wine Yeasts*. Research Signpost, Kerala, India, pp. 35–52.
- Gognies, S., Barka, E.A., Gainvors-Claiss, A., Belarbi, A., 2006. Interactions between yeasts and grapevines: filamentous growth, endopolygalacturonase and phytopathogenicity of colonizing yeasts. *Microbial Ecology* 51, 109–116.
- Golubev, W.I., 2006. Antagonistic interactions among yeasts. In: Rosa, C.A., Péter, G. (Eds.), *The Yeast Handbook. Biodiversity and Ecophysiology of Yeasts*, Springer, Berlin, Germany, pp. 197–219.
- Gomes, A., Sousa, S., Gomes, P., Santos, B., Carreto, L., Santos, S., Egas, C., Santos, A., 2009. Assessment of Microbial Diversity in Wine Musts. Proceedings of Microsafetywine Symposium, Villafranca del Penedès, Spain.
- González, Á., Hierro, N., Poblet, M., Rozés, N., Mas, A., Guillamón, J.M., 2004. Application of molecular methods for the differentiation of acetic acid bacteria in a red wine fermentation. *Journal of Applied Microbiology* 96, 853–860.
- González, Á., Hierro, N., Poblet, M., Mas, A., Guillamón, J.M., 2005. Application of molecular methods to demonstrate species and strain evolution of acetic acid bacteria population during wine production. *International Journal of Food Microbiology* 102, 295–304.
- González, Á., Guillamón, J.M., Mas, A., Poblet, M., 2006a. Application of molecular methods for routine identification of acetic acid bacteria. *International Journal of Food Microbiology* 108, 141–146.
- González, Á., Hierro, N., Poblet, M., Mas, A., Guillamón, J.M., 2006b. Enumeration and detection of acetic acid bacteria by real-time PCR and nested PCR. *FEMS Microbiology Letters* 254, 123–128.
- Groenewald, W.H., van Reenen, C.A., Dicks, L.M.T., 2006. Strains of *Lactobacillus plantarum* in grape must are also present in the intestinal tract of vinegar flies. *South African Journal for Enology and Viticulture* 27, 46–50.
- Guerra, E., Sordi, G., Mannazu, I., Clementi, F., Faticenti, F., 1999. Occurrence of wine yeasts on grapes subjected to different pesticide treatments. *Italian Journal of Food Science* 11, 221–230.
- Guerzoni, M.E., Marchetti, R., 1982. Microflora associata al marciume acido della vite e modificazioni indotte dalla malattia sulla composizione di uve e mosti. *Difesa Piante* 4, 231–246.
- Guerzoni, E., Marchetti, R., 1987. Analysis of yeast flora associated with grape sour rot and of the chemical disease markers. *Applied and Environmental Microbiology* 53, 571–576.
- Guillamón, J.M., Mas, A., 2011. Acetic acid bacteria. In: Carrascosa, A.V., Muñoz, R., González, R. (Eds.), *Molecular Wine Microbiology*, 1st ed. Elsevier, Academic Press, London, UK, pp. 227–255.
- Guillamón, J.M., Sabaté, J., Barrio, E., Cano, J., Querol, A., 1998. Rapid identification of wine yeast species based on RFLP analysis of the ribosomal internal transcribed spacer (ITS) region. *Archives of Microbiology* 169, 387–392.
- Gullo, M., Caggia, C., De Vero, L., Giudici, P., 2006. Characterization of acetic acid bacteria in “traditional balsamic vinegar”. *International Journal of Food Microbiology* 106, 209–212.
- Gullo, M., De Vero, L., Giudici, P., 2009. Succession of selected strains of *acetobacter pasteurianus* and other acetic acid bacteria in traditional balsamic vinegar. *Applied and Environmental Microbiology* 75, 2585–2589.
- Huson, D., Auch, A.F., Qi, J., Schuster, S.C., 2007. MEGAN analysis of metagenomic data. *Genome Research* 17, 377–386.
- Illaca, C., Navarrete, P., Mardones, P., Romero, J., Mas, A., 2008. Application of culture independent molecular biology based methods to evaluate acetic acid bacteria diversity during vinegar processing. *International Journal of Food Microbiology* 126, 245–249.
- James, S.A., Cai, J., Roberts, I.N., Collins, M.D., 1997. A phylogenetic analysis of the genus *saccharomyces* based on 18S rRNA gene sequences: description of *saccharomyces kunashirensis* sp. nov. and *saccharomyces martiniae* sp. nov. *International Journal of Systematic Bacteriology* 47, 453–460.

- Jolly, N., Augustyn, O., Pretorius, I., 2003. The occurrence of non-*Saccharomyces cerevisiae* yeast species over three vintages in four vineyards and grape musts from four production regions of the Western Cape, South Africa. *South African Journal of Enology and Viticulture* 24, 35–42.
- Joyeux, A., Lafon-Lafourcade, S., Ribereau-Gayon, P., 1984. Evolution of acetic acid bacteria during fermentation and storage of wine. *Applied and Environmental Microbiology* 48, 153–156.
- Kennelly, M.M., Eugster, C., Gadoury, D.M., Smart, C.D., Seem, R.C., Gobbin, D., Gessler, C., 2005. Contributions of oospore inoculum to epidemics of grapevine downy mildew (*Plasmopara viticola*). *Phytopathology* 94, S50.
- Khan, W., Augustyn, O.P.H., van der Westhuizen, T.J., Lambrechts, M.G., Pretorius, I.S., 2000. Geographic distribution and evaluation of *Saccharomyces cerevisiae* strains isolated from vineyards in the warmer, inland regions of the Western Cape in South Africa. *South African Journal of Enology and Viticulture* 21, 17–31.
- Kish, S., Sharf, R., Margalith, P., 1983. A note on a selective medium for wine yeasts. *Journal of Applied Microbiology* 55, 177–179.
- Kreger-van Rij, N.J.W., 1984. *The Yeast, A Taxonomic Study*, 3rd ed. Elsevier Science B.V., Amsterdam, the Netherlands.
- Krieg, N.R., Holt, J.G. (Eds.), 1984–1989. *Bergey's Manual of Systematic Bacteriology*, vol. 1–4. Williams & Wilkins, Baltimore.
- Kurtzman, C.P., Fell, J.W., Boekhout, T. (Eds.), 2011. *The Yeasts, A Taxonomic Study*, 5th edn. Elsevier, Amsterdam.
- Kurtzman, C.P., Robnett, C.J., 1998. Identification and phylogeny of ascomycetous yeasts from analysis of nuclear large subunit (26S) ribosomal DNA partial sequences. *Antonie van Leeuwenhoek* 73, 331–371.
- La Guerche, S., Dauphin, B., Pons, M., Blancard, D., Darriet, P., 2006. Characterization of some mushroom and earthy off-odors microbially induced by the development of rot on grapes. *Journal of Agricultural and Food Chemistry* 54, 9193–9200.
- Lachance, M.-A., 2003. The Phaff school of yeast ecology. *International Microbiology* 6, 163–167.
- Lafon-Lafourcade, S., Carre, E., Ribereau-Gayon, P., 1983. Occurrence of lactic acid bacteria during the different stages of vinification and conservation of wines. *Applied and Environmental Microbiology* 46, 874–880.
- Leong, S.-L., Hocking, A.D., Scott, E.S., 2006. Survival and growth of *Aspergillus carbonarius* on wine grapes before harvest. *International Journal of Food Microbiology* 111 (Supplement 1), S83–S87.
- Lerm, E., Engelbrecht, L., du Toit, M., 2010. Malolactic Fermentation: The ABC's of MLF. *South African Journal of Enology and Viticulture* 31, 186–212.
- Li, S.-S., Cheng, C., Li, Z., Chen, J.-Y., Yan, B., Han, B.-Z., Reeves, M., 2010. Yeast species associated with wine grapes in China. *International Journal of Food Microbiology* 138, 85–90.
- Logrieco, A., Ferracane, R., Cozzi, G., Haidukowsky, M., Susca, A., Mulè, G., Ritieni, A., 2011. Fumonisin B2 by *Aspergillus niger* in the grape-wine chain: an additional potential mycotoxicological risk. *Annals of Microbiology* 61, 1–3.
- Loinger, C., Cohen, S., Dror, N., Berlinger, M.J., 1977. Effect of grape cluster rot on wine quality. *American Journal of Enology and Viticulture* 28, 196–199.
- Longo, E., Cansado, J., Agrelo, D., Villa, T.G., 1991. Effect of climatic conditions on yeast diversity in grape musts from Northwest Spain. *Am. J. Enol. Vitic.* 42, 141–144.
- Lopez, I., Ruiz-Larrea, F., Cocolin, L., Orr, E., Phister, T., Marshall, M., VanderGheynst, J., Mills, D.A., 2003. Design and evaluation of PCR primers for analysis of bacterial populations in wine by denaturing gradient gel electrophoresis. *Applied and Environmental Microbiology* 69, 6801–6807.
- Loureiro, V., Malfeito-Ferreira, M., 2003. Spoilage yeasts in the wine industry. *International Journal of Food Microbiology* 86, 23–50.
- Loureiro, V., Malfeito-Ferreira, M., Carreira, A., 2004. Detecting spoilage yeasts. In: Steele, R. (Ed.), *Understanding and Measuring the Shelf-life of Food*. Woodhead Publishers, Cambridge, pp. 233–288.
- Loureiro, V., Malfeito-Ferreira, M., Monteiro, S., Ferreira, R.B., in press. The microbial community of grape berry. In: Gerós, H., Chaves, M.M., Delrot, S. (Eds.), *The Biochemistry of the Grape Berry*. Bentham Science Publishers.
- Magyar, I., Bene, Z., 2006. Morphological and taxonomic study on mycobiota of noble rotted grapes in the Tokaj wine district. *Acta Alimentaria* 35, 237–246.
- Malfeito-Ferreira, M., 2011. Yeasts and wine off-flavours: a technological perspective. *Annals of Microbiology* 61, 95–102.
- Malfeito-Ferreira, M., Barata, A., Loureiro, V., 2009. Wine spoilage by fungal metabolites. In: Moreno-Arribas, M., Polo, M. (Eds.), *Wine Chemistry and Biochemistry*. Springer, New York, pp. 615–645.
- Marchetti, R., Guerzoni, E., Gentile, M., 1984. Research on the etiology of a new disease of grapes: sour rot. *Vitis* 23, 55–65.
- Martini, A., 1993. Origin and domestication of the wine yeast *Saccharomyces cerevisiae*. *Journal of Wine Research* 4, 165–176.
- Martini, A., Federici, F., Rosini, G., 1980. A new approach to the study of yeast ecology of natural substrates. *Canadian Journal of Microbiology* 26, 856–859.
- Meunier, M., Steel, C.C., 2009. Effect of *Colletotrichum acutatum* ripe rot on the composition and sensory attributes of Cabernet Sauvignon grapes and wine. *Australian Journal of Grape and Wine Research* 15, 223–227.
- Miller, M.W., Phaff, H.J., 1962. Successive microbial populations in Calimyrna figs. *Applied and Environmental Microbiology* 10, 394–400.
- Mills, D.A., Johannsen, E.A., Cocolin, L., 2002. Yeast diversity and persistence in *Botrytis*-affected wine fermentations. *Applied and Environmental Microbiology* 68, 4884–4893.
- Moore, K.J., Johnson, M.G., Morris, J.R., 1988. Indigenous yeast microflora on Arkansas white Riesling (*Vitis-vinifera*) grapes and in model must systems. *Journal of Food Science* 53, 1725–1728.
- Mortimer, R., Polsinelli, M., 1999. On the origins of wine yeast. *Research in Microbiology* 150, 199–204.
- Munöz, R., Moreno-Arribas, M., de las Rivas, B., 2011. Lactic acid bacteria. In: Carrascosa, A.V., Muñoz, R., González, R. (Eds.), *Molecular Wine Microbiology*, 1st ed. Elsevier, Academic Press, London, UK, pp. 191–226.
- Muyzer, G., de Waal, E.C., Uitterlinden, A.G., 1993. Profiling of complex microbial populations by denaturing gradient gel electrophoresis analysis of polymerase chain reaction-amplified genes coding for 16S rRNA. *Applied and Environmental Microbiology* 59, 695–700.
- Nisiotou, A.A., Nychas, G.J.E., 2007. Yeast populations residing on healthy or *Botrytis*-infected grapes from a vineyard in Attica, Greece. *Applied and Environmental Microbiology* 73, 2765–2768.
- Nisiotou, A.A., Rantsiou, K., Iliopoulos, V., Cocolin, L., Nychas, G.-J.E., 2011. Bacterial species associated with sound and *Botrytis*-infected grapes from a Greek vineyard. *International Journal of Food Microbiology* 145, 432–436.
- Noble, A.C., 1998. Why do wines taste bitter and feel astringent? In: Ebeler, S.E., Waterhouse, A.L. (Eds.), *Chemistry of Wine Flavor*. ACS Symposium Series 714. American Chemical Society, Washington, D.C. pp. 156–165.
- Oliva, J., Navarro, S., Navarro, G., Cámara, M.A., Barba, A., 1999. Integrated control of grape berry moth (*Lobesia botrana*), powdery mildew (*Uncinula necator*), downy mildew (*Plasmopara viticola*) and grapevine sour rot (*Acetobacter* spp.). *Crop Protection* 18, 581–587.
- Oliva, J., Cayuela, M., Paya, P., Martínez-Cacha, A., Cámara, M.A., Barba, A., 2007. Influence of fungicides on grape yeast content and its evolution in the fermentation. *Communications in Agricultural and Applied Biological Sciences* 72, 181–189.
- Pan, W., Jussier, D., Terrade, N., Yada, R.Y., Mira de Orduña, R., 2011. Kinetics of sugars, organic acids and acetaldehyde during simultaneous yeast-bacterial fermentations of white wine at different pH values. *Food Research International* 44, 660–666.
- Poblet, M., Rozès, N., Guillamón, J.M., Mas, A., 2000. Identification of acetic acid bacteria by restriction fragment length polymorphism analysis of a PCR-amplified fragment of the gene coding for 16S rRNA. *Letters in Applied Microbiology* 31, 63–67.
- Pons, A., Lavigne, V., Eric, F., Darriet, P., Dubourdieu, D., 2008. Identification of volatile compounds responsible for prune aroma in prematurely aged red wines. *Journal of Agricultural and Food Chemistry* 56, 5285–5290.
- Poulard, A., Simon, L., Cuinier, C., 1981. Caractères de la microflore levurienne du vignoble Nantais. *Vignes Vins* 300, 14–18.
- Prakitchaiwattana, C., 2004. Investigation of Yeasts Associated with Australian Wine Grapes using Cultural and Molecular Methods. University of New South Wales, PhD Thesis.
- Prakitchaiwattana, C.J., Fleet, G.H., Heard, G.M., 2004. Application and evaluation of denaturing gradient gel electrophoresis to analyse the yeast ecology of wine grapes. *Fems Yeast Research* 4, 865–877.
- Pretorius, I.S., Van der Westhuizen, T.J., Augustyn, O.P.H., 1999. Yeast Biodiversity in Vineyards and Wineries and Its Importance to the South African Wine Industry. A Review. *South African Journal of Enology and Viticulture* 20.
- Prieto, C., Jara, C., Mas, A., Romero, J., 2007. Application of molecular methods for analysing the distribution and diversity of acetic acid bacteria in Chilean vineyards. *International Journal of Food Microbiology* 115, 348–355.
- Querol, A., Fleet, G. (Eds.), 2006. *Yeasts in Food and Beverages*. Springer, Berlin, Heidelberg.
- Rao, R.S., Bhadra, B., Kumar, N.N., Shivaji, S., 2007. *Candida hyderabadensis* sp. nov., a novel ascomycetous yeast isolated from wine grapes. *Fems Yeast Research* 7, 489–493.
- Raspor, P., Milek, D.M., Polanc, J., Smole Mozina, S., Cadez, N., 2006. Yeasts isolated from three varieties of grapes cultivated in different locations of the Dolenjska vinegrowing region, Slovenia. *International Journal of Food Microbiology* 109, 97–102.
- Regueiro, L.A., Costas, C.L., Rubio, J.E.L., 1993. Influence of viticultural and enological practices on the development of yeast populations during winemaking. *American Journal of Enology and Viticulture* 44, 405–408.
- Rementeria, A., Rodriguez, J.A., Cadaval, A., Amenabar, R., Muguruza, J.R., Hernando, F.L., Sevilla, M.J., 2003. Yeast associated with spontaneous fermentations of white wines from the "Txakoli de Bizkaia" region (Basque Country, North Spain). *International Journal of Food Microbiology* 86, 201–207.
- Renouf, V., Lonvaud-Funel, A., 2007. Development of an enrichment medium to detect *Dekkera/Brettanomyces bruxellensis*, a spoilage wine yeast, on the surface of grape berries. *Microbiological Research* 162, 154–167.
- Renouf, V., Claisse, O., Lonvaud-Funel, A., 2005. Understanding the microbial ecosystem on the grape berry surface through numeration and identification of yeast and bacteria. *Australian Journal of Grape and Wine Research* 11, 316–327.
- Renouf, V., Claisse, O., Lonvaud-Funel, A., 2007. Inventory and monitoring of wine microbial consortia. *Applied Microbiology and Biotechnology* 75, 149–164.
- Rodas, A.M., Ferrer, S., Pardo, I., 2003. 16S-ARDRA, a tool for identification of lactic acid bacteria isolated from grape must and wine. *Systematic and Applied Microbiology* 26, 412–422.
- Rodas, A.M., Ferrer, S., Pardo, I., 2005. Polyphasic study of wine *Lactobacillus* strains: taxonomic implications. *International Journal of Systematic and Evolutionary Microbiology* 55, 197–207.
- Rodrigues, F., Goncalves, G., Pereira-da-Silva, S., Malfeito-Ferreira, M., Loureiro, V., 2001. Development and use of a new medium to detect yeasts of the genera *Dekkera/Brettanomyces*. *Journal of Applied Microbiology* 90, 588–599.
- Rogiers, S.Y., Whitelaw-Weckert, M., Radovanovic-Tesic, M., Greer, L.A., White, R.G., Steel, C.C., 2005. Effects of spray adjuvants on grape (*Vitis vinifera*) berry microflora, epicuticular wax and susceptibility to infection by *Botrytis cinerea*. *Australasian Plant Pathology* 34, 221–228.
- Romano, P., 2002. Role of apiculate yeasts on organoleptic characteristics of wine. In: Ciani, M. (Ed.), *Biodiversity and Biotechnology of Wine Yeasts*. Research Signpost, Kerala, India, pp. 99–109.
- Rosa, C., Péter, G. (Eds.), 2006. *Biodiversity and Ecophysiology of Yeasts*. Springer Berlin, Heidelberg.

- Ruiz, A., Poblet, M., Mas, A., Guillamon, J., 2000. Identification of acetic acid bacteria by RFLP of PCR-amplified 16S rDNA and 16S–23S rDNA intergenic spacer. *International Journal of Systematic and Evolutionary Microbiology* 50, 1981–1987.
- Sabate, J., Cano, J., Esteve-Zarzoso, B., Guillamón, J.M., 2002. Isolation and identification of yeasts associated with vineyard and winery by RFLP analysis of ribosomal genes and mitochondrial DNA. *Microbiological Research* 157, 267–274.
- Sampaio, J.P., Gonçalves, P., 2008. Natural populations of *Saccharomyces kudriavzevii* in Portugal are associated with Oak bark and are sympatric with *S. cerevisiae* and *S. paradoxus*. *Applied and Environmental Microbiology* 74, 2144–2152.
- Sangorrín, M.P., Zajonskovsky, I.E., Lopes, C.A., Rodríguez, M.E., Giraudo de van Broock, M.R., Caballero, A.C., 2001. Killer behaviour in wild wine yeasts associated with Merlot and Malbec type musts spontaneously fermented from Northwestern Patagonia (Argentina). *Journal of Basic Microbiology* 41, 105–113.
- Santos, C., Fraga, M.E., Kozakiewicz, Z., Lima, N., 2010. Fourier transform infrared as a powerful technique for the identification and characterization of filamentous fungi and yeasts. *Research in Microbiology* 161, 168–175.
- Sarrazin, E., Dubourdieu, D., Darriet, P., 2007. Characterization of key-aroma compounds of botrytized wines, influence of grape botrytization. *Food Chemistry* 103, 536–545.
- Schuller, D., Casal, M., 2007. The genetic structure of fermentative vineyard-associated *Saccharomyces cerevisiae* populations revealed by microsatellite analysis. *Antonie van Leeuwenhoek* 91, 137–150.
- Schuller, D., Córte-Real, M., Leão, C., 2000. A differential medium for the enumeration of the spoilage yeast *Zygosaccharomyces bailii* in wine. *Journal of Food Protection* 63, 1570–1575.
- Schuller, D., Alves, H., Dequin, S., Casal, M., 2005. Ecological survey of *saccharomyces cerevisiae* strains from vineyards in the Vinho Verde region of Portugal. *FEMS Microbiology Ecology* 51, 167–177.
- Serra, R., Peterson, S.W., 2007. *Penicillium astrolabium* and *Penicillium neocrassum*, two new species isolated from grapes and their phylogenetic placement in the *P. olsonii* and *P. brevicompactum* clade. *Mycologia* 99, 78–87.
- Sigrist, I.A., Manzardo, G., Amadô, R., 2001. Analysis of furan fatty acids in dried green herbs using ion trap GC–MS/MS. In: Pfannhauser, W., Fenwick, G.R., Khokhar, S. (Eds.), *Biologically-active Phytochemicals in Food*. The Royal Society of Chemistry, Cambridge, UK, pp. 237–240.
- Sipiczki, M., 2003. *Candida zemplinina* sp. nov., an osmotolerant and psychrotolerant yeast that ferments sweet botrytized wines. *International Journal of Systematic and Evolutionary Microbiology* 53, 2079–2083.
- Skelin, A., Sikora, S., Orlić, S., Duraković, L., Redžepović, S., 2008. Genetic diversity of indigenous *Saccharomyces sensu stricto* yeasts isolated from southern Croatia. *Agri-culturae Conspectus Scientificus* 73, 89–94.
- Steel, C.C., Greer, L.A., Savocchia, S., 2007. Studies on *Colletotrichum acutatum* and *Greeneria uvicola*: two fungi associated with bunch rot of grapes in sub-tropical Australia. *Australian Journal of Grape and Wine Research* 13, 23–29.
- Steel, C.C., Greer, L.A., Savocchia, S., Samuelian, S.K., 2011. Effect of temperature on *Botrytis cinerea*, *Colletotrichum acutatum* and *Greeneria uvicola* mixed fungal infection of *Vitis vinifera* grape berries. *Vitis: Journal of Grapevine Research* 50, 69–71.
- Stevic, S., 1962. The significance of bees (*Apis* sp.) and wasps (*Vespa* sp.) as carriers of yeast for the microflora of grapes and the quality of wine. *Arkh. Poljoprivredne Nauke* 50, 80–92.
- Stummer, B.E., Francis, I.L., Zanker, T., Lattey, K.A., Scott, E.S., 2005. Effects of powdery mildew on the sensory properties and composition of Chardonnay juice and wine when grape sugar ripeness is standardised. *Australian Journal of Grape and Wine Research* 11, 66–76.
- Sturm, J., Grossmann, M., Schnell, S., 2006. Influence of grape treatment on the wine yeast populations isolated from spontaneous fermentations. *Journal of Applied Microbiology* 101, 1241–1248.
- Subden, R.E., Husnik, J.I., van Twest, R., van der Merwe, G., van Vuuren, H.J.J., 2003. Autochthonous microbial population in a Niagara Peninsula icewine must. *Food Research International* 36, 747–751.
- Tat, L., Comuzzo, P., Battistutta, F., Zironi, R., 2007. Sweet-like off-flavor in Aglianico del Vulture wine: ethyl phenylacetate as the mainly involved compound. *Journal of Agricultural and Food Chemistry* 55, 5205–5212.
- Thibon, C., Dubourdieu, D., Darriet, P., Tominaga, T., 2009. Impact of noble rot on the aroma precursor of 3-sulfanylhexanol content in *Vitis vinifera* L. cv Sauvignon blanc and Semillon grape juice. *Food Chemistry* 114, 1359–1364.
- Torija, M.J., Mateo, E., Guillamón, J.M., Mas, A., 2010. Identification and quantification of acetic acid bacteria in wine and vinegar by TaqMan-MGB probes. *Food Microbiology* 27, 257–265.
- Urso, R., Rantsiou, K., Dolci, P., Rolle, L., Comi, G., Cocolin, L., 2008. Yeast biodiversity and dynamics during sweet wine production as determined by molecular methods. *Fems Yeast Research* 8, 1053–1062.
- Valero, E., Schuller, D., Cambon, B., Casal, M., Dequin, S., 2005. Dissemination and survival of commercial wine yeast in the vineyard: a large-scale, three-year study. *Fems Yeast Research* 5, 959–969.
- Valero, E., Cambon, B., Schuller, D., Casal, M., Dequin, S., 2007. Biodiversity of *Saccharomyces* yeast strains from grape berries of wine-producing areas using starter commercial yeasts. *Fems Yeast Research* 7, 317–329.
- Van der Westhuizen, T.J., Augustyn, O.P.H., Khan, W., Pretorius, I.S., 2000a. Seasonal Variation of Indigenous *Saccharomyces cerevisiae* Strains Isolated from Vineyards of the Western Cape in South Africa. *South African Journal of Enology and Viticulture* 21, 10–16.
- van der Westhuizen, T.J., Augustyn, O.P.H., Pretorius, I.S., 2000b. Geographical distribution of indigenous *Saccharomyces cerevisiae* strains isolated from vineyards in the coastal regions of the Western Cape in South Africa. *South African Journal for Enology and Viticulture* 21, 3–9.
- Vaughan-Martini, A., Martini, A., 1995. Facts, myths and legends on the prime industrial microorganism. *Journal of Industrial Microbiology* 14, 514–522.
- Velišek, J., Cejpek, K., 2006. Biosynthesis of food constituents: lipids. 1. Fatty acids and derived compounds — a review. *Czech Journal of Food Sciences* 24, 193–216.
- Ventura, M., Casas, I.A., Morelli, L., Callegari, M.L., 2000. Rapid amplified ribosomal DNA restriction analysis (ARDRA) identification of *Lactobacillus* spp. isolated from fecal and vaginal samples. *Systematic and Applied Microbiology* 23, 504–509.
- Viviani-Nauer, A., Hoffmann-Boller, P., Basler, P., Gafner, J., 1995. Wild yeast flora on grapes of fungi disease resistant cultivars and their dynamics during fermentation. *Schweizerische Zeitschrift fuer Obst-und Weinbau, Waedenswil* 131, 390–393.
- Wenning, M., Seiler, H., Scherer, S., 2002. Fourier-Transform Infrared Microspectroscopy, a Novel and Rapid Tool for Identification of Yeasts. *Applied and Environmental Microbiology* 68, 4717–4721.
- Wenning, M., Büchl, N.R., Scherer, S., 2010. Species and strain identification of lactic acid bacteria using FTIR spectroscopy and artificial neural networks. *Journal of Biophotonics* 3, 493–505.
- Yamada, Y., Yukphan, P., 2008. Genera and species in acetic acid bacteria. *International Journal of Food Microbiology* 125, 15–24.
- Yanagida, F., Ichinose, F., Shinohara, T., Goto, S., 1992. Distribution of wild yeasts in the white grape varieties at Central Japan. *The Journal of General and Applied Microbiology* 38, 505–509.
- Zahavi, T., Drobny, S., Cohen, L., Weiss, B., Ben-Arie, R., 2002. Characterization of the yeast flora on the surface of grape berries in Israel. *Vitis: Journal of Grapevine Research* 41, 203–208.