



# Article Primary Metabolite Profile Changes in *Coffea* spp. Promoted by Single and Combined Exposure to Drought and Elevated CO<sub>2</sub> Concentration

Ana M. Rodrigues <sup>1,+</sup><sup>(D)</sup>, Tiago Jorge <sup>1</sup>, Sonia Osorio <sup>2</sup><sup>(D)</sup>, Delphine M. Pott <sup>2</sup><sup>(D)</sup>, Fernando C. Lidon <sup>3</sup>, Fábio M. DaMatta <sup>4</sup><sup>(D)</sup>, Isabel Marques <sup>5</sup><sup>(D)</sup>, Ana I. Ribeiro-Barros <sup>3,5,\*</sup><sup>(D)</sup>, José C. Ramalho <sup>3,5,\*</sup><sup>(D)</sup> and Carla António <sup>1,\*,+</sup><sup>(D)</sup>

- <sup>1</sup> Plant Metabolomics Laboratory, Instituto de Tecnologia Química e Biológica António Xavier, Universidade Nova de Lisboa (ITQB NOVA), 2780-157 Oeiras, Portugal; amrodrigues@itqb.unl.pt (A.M.R.); tiago.jorge89@gmail.com (T.J.)
- <sup>2</sup> Instituto de Hortofruticultura Subtropical y Mediterránea "La Mayora", Departamento de Biología Molecular y Bioquímica, Universidad de Málaga—Consejo Superior de Investigaciones Científicas (IHSM-UMA-CSIC), 29071 Málaga, Spain; sosorio@uma.es (S.O.); dpott@uma.es (D.M.P.)
- <sup>3</sup> GeoBioSciences, GeoTechnologies and GeoEngineering (GeoBioTec), Faculdade de Ciências e Tecnologia (FCT), Universidade NOVA de Lisboa (UNL), 2829-516 Monte de Caparica, Portugal; fjl@fct.unl.pt
- <sup>4</sup> Departamento de Biologia Vegetal, Universidade Federal Viçosa (UFV), Viçosa 36570-090, Brazil; fdamatta@ufv.br
- <sup>5</sup> Plant Stress & Biodiversity Lab, Centro de Estudos Florestais (CEF), Instituto Superior Agronomia (ISA), Universidade de Lisboa (ULisboa), Tapada da Ajuda, 1349-017 Lisboa, Portugal; isabelmarques@isa.ulisboa.pt
- \* Correspondence: aribeiro@isa.ulisboa.pt (A.I.R.-B.); cochichor@mail.telepac.pt (J.C.R.);
  - plantmetabolomicslabpt@gmail.com (C.A.)
  - Current address: Plant Metabolomics Lab Portugal, Centro de Estudos Florestais (CEF), Instituto Superior Agronomia (ISA), Universidade de Lisboa (ULisboa), Tapada da Ajuda, 1349-017 Lisboa, Portugal.

**Abstract:** Climate change scenarios pose major threats to many crops worldwide, including coffee. We explored the primary metabolite responses in two *Coffea* genotypes, *C. canephora* cv. Conilon Clone 153 and *C. arabica* cv. Icatu, grown at normal (aCO<sub>2</sub>) or elevated (eCO<sub>2</sub>) CO<sub>2</sub> concentrations of 380 or 700 ppm, respectively, under well-watered (WW), moderate (MWD), or severe (SWD) water deficit conditions, in order to assess coffee responses to drought and how eCO<sub>2</sub> can influence such responses. Primary metabolites were analyzed with a gas chromatography time-of-flight mass spectrometry metabolomics platform (GC-TOF-MS). A total of 48 primary metabolites were identified in both genotypes (23 amino acids and derivatives, 10 organic acids, 11 sugars, and 4 other metabolites), with differences recorded in both genotypes. Increased metabolite levels were observed in CL153 plants under single and combined conditions of aCO<sub>2</sub> and drought (MWD and SWD), as opposed to the observed decreased levels under eCO<sub>2</sub> in both drought conditions. In contrast, Icatu showed minor differences under MWD, and increased levels (especially amino acids) only under SWD at both CO<sub>2</sub> concentration conditions, although with a tendency towards greater increases under eCO<sub>2</sub> in either MWD and SWD, in contrast with Icatu.

**Keywords:** climate change; *Coffea arabica; Coffea canephora;* coffee tree; elevated CO<sub>2</sub>; GC-TOF-MS; mass spectrometry; plant metabolomics; water deficit

# 1. Introduction

Coffee is an important agricultural commodity worldwide that provides income and employment to millions of people in developing countries. Coffee is cultivated in over 80 countries throughout the tropics with ca. 99% of the production based on *Coffea arabica* L. (Arabica coffee) and *Coffea canephora* Pierre ex Froehner (Robusta coffee) [1,2]. Estimated global climate change scenarios will impose adverse environmental conditions with deep



Citation: Rodrigues, A.M.; Jorge, T.; Osorio, S.; Pott, D.M.; Lidon, F.C.; DaMatta, F.M.; Marques, I.; Ribeiro-Barros, A.I.; Ramalho, J.C.; António, C. Primary Metabolite Profile Changes in *Coffea* spp. Promoted by Single and Combined Exposure to Drought and Elevated CO<sub>2</sub> Concentration. *Metabolites* **2021**, *11*, 427. https:// doi.org/10.3390/metabol1070427

Academic Editor: Hirokazu Kawagishi

Received: 17 June 2021 Accepted: 25 June 2021 Published: 29 June 2021

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https:// creativecommons.org/licenses/by/ 4.0/). impacts on crops, mainly due to water constraints and heat stress [3]. In this context, the coffee sector is expected to face serious challenges in the upcoming decades as several studies have demonstrated the climate change sensitivity of coffee species, particularly *C. arabica*, with impacts on suitable cultivation areas, yield, and bean quality, as well as increased pest and disease incidence and economic losses [4–9].

One of the major components of global climate change is an increase in air CO<sub>2</sub> concentration, coupled with rising air temperatures (with the worst estimate pointing to a global average increase of up to 936 ppm  $CO_2$  and 4.5 °C by 2100), together with more frequent and prolonged drought events associated with unpredictable rainfall patterns [10]. Such events are expected to threaten the sustainability of agricultural production on a global scale and be responsible for severe constraints to plant growth and productivity [11]. Nevertheless, there are interesting and potentially positive effects of elevated CO<sub>2</sub> concentration (eCO<sub>2</sub>) on several crop plants, usually associated with increased photosynthetic activity [12,13]. That is the case for coffee, under adequate water availability [14–17], which displays greater C-investment in reproductive structures [18], ultimately increasing productivity [1]. Furthermore, eCO<sub>2</sub> was recently reported to increase coffee plant resilience to heat stress [19–21], while preserving leaf mineral balance [22] and bean quality [23]. Additionally, although the future positive impact of eCO<sub>2</sub> on plants subject to water deficits can be species-dependent [24], studies reported that  $eCO_2$  can clearly reduce drought impact on plant photosynthesis, growth, and yields [25–27]). These effects were reported to occur in *Coffea* spp. as well, where  $eCO_2$  mitigated the negative impacts of drought on photosynthetic performance and components [28–32].

Under field conditions, plants are often exposed to a combination of different environmental changes. In this case, the plant response to combined changing conditions should be regarded not as a sum of each one applied independently, but as a unique response that can reflect a positive (i.e., increased tolerance) or negative (i.e., susceptibility) interaction between them [33–35]. Abiotic stresses can deeply impact plant metabolism and cause a reconfiguration of the metabolic network in order to allow for the maintenance of metabolic homeostasis and the production of compounds that can mitigate the effects of stress [36,37]. Drought stress is one of the major threats to agriculture, since it constrains a wide number of physiological, morphological, and biochemical processes, impacting growth, nutrient uptake, C-assimilation, and partitioning, and ultimately yield and quality [28,38,39]. Hence, it is essential to understand the plant response to this critical adverse environmental condition. Metabolic adjustments in response to drought include the regulation of photosynthesis and the maintenance of cell osmotic potential through the accumulation of metabolites such as phytohormone abscisic acid and osmolytes, including amino acids, soluble sugars, the raffinose family of oligosaccharides (RFOs), polyols, and polyamines [36,40–43]. Osmolyte accumulation in response to abiotic stress conditions has an important role in the maintenance of cell turgor—via a decrease in the osmotic potential of the cell—and protection against oxidative damage by a decrease in the levels of reactive oxygen species (ROS) to restore cellular redox balance [43]. Moreover, stress-tolerant plants usually show higher levels of some of these stress-related metabolites, even under normal growth conditions, to keep their metabolism prepared for adverse environmental conditions [41].

Complementing the previous studies of our team, and regarding the coffee plant response to single and combined exposure to drought and/or eCO<sub>2</sub> at the physiological, biochemical, and molecular level [28,29,32,44], here we further explored the changes in gas chromatography time-of-flight mass spectrometry (GC-TOF-MS) primary metabolite profiling in plants exposed to increasing water deficit severity, and how eCO<sub>2</sub> might influence drought response and plant resilience in *Coffea* spp. To date, GC-MS is the most widely accepted analytical technology used in post-genomic plant metabolomics studies [43,45–47]. Genotypes from the two main coffee producing species, *C. canephora* cv. Conilon Clone 153 (CL153) and *C. arabica* cv. Icatu (Icatu), were grown under normal atmospheric (aCO<sub>2</sub>) or elevated (eCO<sub>2</sub>) air CO<sub>2</sub> concentration, coupled with either well-

watered conditions (WW), or moderate (MWD) or severe (SWD) water deficit conditions. Our findings provide important and timely evidence regarding the role of  $eCO_2$  in the response to drought in these species, contributing to improvements in knowledge regarding coffee plant performance in the face of ongoing and future global climate changes.

### 2. Results

The GC-TOF-MS analysis identified 48 primary metabolites in CL153 and Icatu leaves, including 23 amino acids and derivatives, 10 organic acids, 11 sugars, and 4 other metabolites (Figures 1 and 2, Supplementary Tables S1 and S2).





three water availability levels (WW, well-watered; MWD, moderate water deficit; SWD, severe water deficit). Results indicate (**a**) single exposure to eCO<sub>2</sub>, (**b**) single exposure to each water deficit level under aCO<sub>2</sub>, (**c**) exposure to each water deficit level under eCO<sub>2</sub>, and (**d**) eCO<sub>2</sub> effect within each water deficit level. Relative values (as means of three to six independent measurements) were normalized to the internal standard (ribitol) and the dry weight of the samples; false color imaging was performed on log<sub>10</sub>-transformed GC-TOF-MS metabolite data. Significant changes using one-way ANOVA are indicated as ( $\bigcirc$ ) for *p* < 0.05. Heatmaps and clustering were performed in R software [48,49] using the "gplots" package [50]. Gray-color squares represent non-detected values. Metabolites are grouped in amino acids and derivatives (AA), organic acids (OA), sugars and sugar alcohols (SS), and others (O).



Figure 2. Heatmap representing the changes in relative levels of primary metabolites in leaves of

*Coffea arabica* cv. Icatu (Icatu) grown under 380 (aCO<sub>2</sub>) or 700 (eCO<sub>2</sub>) ppm CO<sub>2</sub>, and three water availability levels (WW, well-watered; MWD, moderate water deficit; SWD, severe water deficit). Results indicate (**a**) single exposure to eCO<sub>2</sub>, (**b**) single exposure to each water deficit level under aCO<sub>2</sub>, (**c**) exposure to each water deficit level under eCO<sub>2</sub>, and (**d**) eCO<sub>2</sub> effect within each water deficit level. Relative values (as means of three to six independent measurements) were normalized to the internal standard (ribitol) and the dry weight of the samples; false color imaging was performed on log<sub>10</sub>-transformed GC-TOF-MS metabolite data. Significant changes using one-way ANOVA are indicated as ( $\bigcirc$ ) for *p* < 0.05. Heatmaps and clustering were performed in R software [48,49] using the "gplots" package [50]. Gray-color squares represent non-detected values. Metabolites are grouped in amino acids and derivatives (AA), organic acids (OA), sugars and sugar alcohols (SS), and others (O).

In general, a larger number of significant changes in primary metabolite levels were observed in CL153 than in Icatu due to single exposure to  $eCO_2$  (WW plants), and MWD (both  $CO_2$  concentrations). However, under SWD, both genotypes showed a large (and similar) number of significant changes under both  $aCO_2$  or  $eCO_2$ . Additionally, in CL153 SWD plants under  $eCO_2$ , those changes reflected mostly a decrease in metabolite levels (as opposed to Icatu) (Figures 1c and 2c). Lower metabolite levels were observed under  $eCO_2$  as compared with  $aCO_2$  in both MWD or SWD (Figure 1d), whereas Icatu revealed opposite patterns under  $eCO_2$ , as compared with  $aCO_2$  (Figure 2d).

In CL153 plants, the effect of single exposure to  $eCO_2$  led to a significant increase in aspartate, glutamate, methionine, pyroglutamate, 2-oxoglutarate, citrate, quinate, and glucose (up to 5-fold) (Figure 1a). Additionally, exposure to single drought conditions caused a higher number of significant changes in primary metabolite levels (Figure 1b). Under MWD, CL153 plants grown under aCO<sub>2</sub> revealed significant increases (up to 5-fold) in most amino acids ( $\gamma$ -aminobutyric acid (GABA), glutamate, histidine, isoleucine, lysine, pyroglutamate, threonine, tryptophan, tyramine, tyrosine, and valine). A similar response was observed under SWD with significant increases in amino acids (up to 11-fold), together with a significant increase in glucose (5-fold), and a significant decline in sucrose, citrate, and phosphoric acid. Interestingly, CL153 plants grown under eCO<sub>2</sub> and subjected to MWD and SWD conditions (Figure 1c) showed a markedly different response than under WW conditions, with reduction of a large number of metabolites. These  $eCO_2$  plants also showed a different pattern under MWD and SWD as compared with their aCO<sub>2</sub> counterparts, revealing significant declines in the levels of asparagine, aspartate, glycine, glutamate (only at SWD), histidine (only at MWD), proline (only at MWD), pyroglutamate, and serine. Nevertheless, under eCO<sub>2</sub>, significant increases in glucose (2-fold, in MWD), and in fructose (4-fold), trehalose (5-fold), tryptophan, (up to 7-fold), and valine (4-fold) were observed under SWD. Significant reductions were observed regarding pyruvate (under MWD), and in citrate, glycerate, and threonate under SWD.

Furthermore, a global reduction in relative metabolite abundance was observed when comparing eCO<sub>2</sub> to aCO<sub>2</sub> for each water deficit treatment in regards to amino acid levels, particularly in MWD (Figure 1d). Still, under the combined exposure to SWD and eCO<sub>2</sub>, only glutamate and pyroglutamate showed a significant decrease, whereas pyruvate and sucrose showed a significant increase (up to 4-fold).

Analysis of *C. arabica* cv. Icatu showed a different pattern of metabolite variation when compared with the *C. canephora* genotype CL153. In Icatu plants, the single eCO<sub>2</sub> effect promoted significant changes in a low number of metabolites (in isoleucine, fructose, and spermine, up to 2-fold) (Figure 2a). Furthermore, in this genotype, only a few metabolites showed altered abundance under MWD, irrespective of the air CO<sub>2</sub> concentration (Figure 2b). Most significant changes were observed in plants under SWD and aCO<sub>2</sub> conditions (Figure 2b,c). Overall, these significant variations included an increase in arginine, fructose, GABA, histidine, isoleucine, lysine, phenylalanine, proline, tryptophan, tyrosine, and valine (up to 9-fold), and a significant decrease in 2-oxoglutarate, aspartate, galactinol, glutamate, glycerate, serine, sucrose, and threonate.

In opposition to what was observed in CL153, comparison between the combined effect of MWD and SWD in plants grown under  $eCO_2$  (Figure 2d) led to few significant changes in primary metabolite levels. These changes included an increase in aspartate (up to 4-fold), pyroglutamate (2-fold, in MWD under  $eCO_2$ ), putrescine, glycerate, threonate (2-fold, in SWD under  $eCO_2$ ), and a significant decrease in arginine and fructose (in MWD under  $eCO_2$ ).

Unsupervised principal component analysis (PCA) was applied to GC-TOF-MS relative primary metabolite data to identify major sources of variation in the data (Figure 3). The CL153 SWD plants clustered separately in relation to WW and MWD plants, whereas a distinction between CO<sub>2</sub> concentration conditions could only be observed for MWD plants.





In Icatu plants (Figure 3b), a clear separation was observed for SWD plants along PC1, in relation to WW and MWD plants. In addition, as observed for CL153 plants, only MWD plants showed a clear distinction between plants grown under aCO<sub>2</sub> and eCO<sub>2</sub>.

To further explore and identify potential stress responsive metabolites in these *Coffea* spp. genotypes, a supervised partial least square discrimination analysis (PLS-DA) was performed (Figure 4). As observed in the PCA, both genotypes showed a clear discrimination between WW and SWD plants, particularly in Icatu plants, regardless of CO<sub>2</sub> concentration. PLS-DA further confirmed a clear distinction between MWD plants grown under different CO<sub>2</sub> concentrations.

The score plots revealed a high number of highly correlated metabolites responsible for the cluster of SWD samples in the negative quadrants of component 1 for both genotypes. These metabolites include mostly amino acids and sugars that significantly increased in plants exposed to single and combined drought and eCO<sub>2</sub> conditions, including the the branched chained amino acids (BCAAs) valine and isoleucine, arginine, GABA, lysine, phenylalanine, trehalose, tryptophan, and tyrosine for CL153 plants, and arginine, BCAAs, fructose, GABA, lysine, proline, tryptophan, and tyrosine for Icatu plants.

var

X-variate 2: 19% expl.

PLS-DA CL153

X-variate 1: 25% expl. var (a)

cle plot CL153

His <sup>Orn</sup> 0.5 0.5 Ery 🔐 Ser TrBro GAC Gł Tym NAc GABA Arg 2 Component 2 roFru Głc Go Component Phe Sen Ph/ Fum\_ Glyc Fri CVE <sup>Arg</sup>√al His DhAsc QNA G3F GABA Euro Pi Cit Lys Put Fum Orn Sp Rhan Erv -0. -0.5 Rai Mal ThrAc NAc Pro Succ<sub>Gły</sub> Asp GAc Se Asp Głu pGlu -1( -0.5 0.5 -0.5 Component 1 Component 1 (d) (c) Figure 4. Partial least square discriminant analysis (PLS-DA) score and correlation circle plots of the primary metabolite profile of leaves of Coffea canephora cv. Conilon Clone 153 (CL153), (a,c), respectively; and of C. arabica cv. Icatu (Icatu), (b,d), respectively, grown under 380 (aCO<sub>2</sub>) or 700 (eCO<sub>2</sub>) ppm CO<sub>2</sub> and increasing water deficit severity; namely, wellwatered (WW), mild water deficit (MWD), and severe water deficit (SWD). Plots performed in R software [48,49] using the

var

X-variate 2: 17% expl.

profile of leaves of *Coffea canephora* cv. Conilon Clone 153 (CL153), (**a**,**c**), respectively; and of *C. arabica* cv. Icatu (Icatu), (**b**,**d**), respectively, grown under 380 (aCO<sub>2</sub>) or 700 (eCO<sub>2</sub>) ppm CO<sub>2</sub> and increasing water deficit severity; namely, well-watered (WW), mild water deficit (MWD), and severe water deficit (SWD). Plots performed in R software [48,49] using the "mixOmics" package [51]. Abbreviations: 2-oxoglutarate (2OG), alanine (Ala), arginine (Arg), asparagine (Asn), aspartate (Asp), citrate (Citr), cystine (cys), dehydroascorbate (DhAsc), erythritol (Ery), fructose (Fru), fucose (Fuc), fumarate (Fum), galactinol (Gol), glucose (Glc), glutamate (Glu), glycerate (GAc), glycerol (Glyc), glycerol-3-phosphate (G3P), glycine (Gly), histidine (His), isoleucine (Ile), lysine (Lys), malate (Mal), methionine (Met), *myo*-Inositol (Ino), nicotinic acid (NAc), ornithine (Orn), phenylalanine (Phe), phosphoric acid (PhA), proline (Pro), putrescine (Put), pyroglutamate (pGlu), pyruvate (Pyr), quinate (QNA), raffinose (Raf), rhamnose (Rham), serine (Ser), spermine (Spm), succinate (Succ), sucrose (Suc), trehalose (Tre), threonate (ThrAc), threonine (Thr), tryptophan (Trp), tyramine (Tym), tyrosine (Tyr), valine (Val).

# 3. Discussion

GC-TOF-MS analysis allowed for the characterization of the primary metabolome of CL153 and Icatu leaves under single and combined exposure to increasing CO<sub>2</sub> concentration (eCO<sub>2</sub>) and water deficit (MWD and SWD) conditions. Unsupervised exploratory PCA revealed differential metabolite responses to single and combined stresses; this effect was clearer in Icatu plants, and PLS-DA analyses allowed us to further identify stress-responsive metabolites whose levels significantly varied upon single and/or combined exposure to drought and eCO<sub>2</sub>.

[CO2] 380 ppm

700 ppm

Water Deficit MWD SWD WW

**PLS-DA** Icatu

X-variate 1: 32% expl. var

Asn

Ino

(b)

Correlation circle plot Icat

pGlu

Thr Met Thra

The significant increase in CL153 alone of most amino acids upon single exposure to MWD indicated that Icatu was not affected by MWD, and thus, supports previous studies reporting its strong drought resilience [32]. However, under SWD, both genotypes showed relevant changes in the levels of most amino acids, although in Icatu these changes were relatively similar under both air CO<sub>2</sub> concentrations. This is also in agreement with the reported drought resilience of this genotype under both CO<sub>2</sub> concentrations [32]. The BCAAs isoleucine and valine significantly increased in Icatu under SWD at both air CO<sub>2</sub> concentrations, and under MWD only at aCO<sub>2</sub>, whereas in CL153 an increase was only observed under SWD and eCO<sub>2</sub>. BCAAs are well-known to accumulate under abiotic stress conditions and play the role of osmolytes during osmotic stress [36,52,53]. BCAAs are also responsible for an alternative supply of electrons and respiratory substrates to the plant respiratory chain under stress conditions, ultimately contributing to drought stress tolerance in plants [54,55].

Lysine, a product of the aspartate pathway, also significantly increased in response to SWD in CL153 plants (only under  $aCO_2$ ), and Icatu (under  $aCO_2$  and  $eCO_2$ ). Lysine catabolism can channel electrons to the mitochondrial electron transport chain in addition to feeding the operation of the tricarboxylic acid (TCA) cycle under carbon starvation [54], as would be the case in these plants due to a strong reduction of photosynthetic C-assimilation under limited water availability [32].

Metabolites from the glutamate family pathway significantly increased in Icatu, including the osmolytes proline and GABA, under SWD irrespective of CO<sub>2</sub> concentration. Osmolyte accumulation has an important role in abiotic stress responses; namely, maintenance of cell turgor via a decrease in the osmotic potential, protection against oxidative damage by decreasing the levels of reactive oxygen species (ROS), and maintenance of the cellular redox balance [41–43]. Accumulation of proline is considered an adaptive mechanism in drought-tolerant coffee genotypes. For instance, C. canephora clones (IC-2, IC-3, IC-4, IC-6, IC-8, and R-4) subjected to drought-stress revealed an accumulation of proline with increasing exposure to soil drying, followed by a decline upon re-watering [56]. Moreover, Silva et al. [57] showed that proline accumulation was higher in the leaves of a droughttolerant clone of *C. canephora* than in a susceptible genotype. Proline levels were also found to increase in C. arabica cvs. Catuaí and Mundo Novo, C. canephora cv. Apoatã, and a graft of Mundo Novo shoot on Apoatã root in response to drought [58]. Arginine, another member of the glutamate family, showed a significant increase in response to the harshest water deficit level under both  $CO_2$  concentrations and in both genotypes. Increased levels of arginine were suggested to contribute to drought tolerance [59]. Arginine is a precursor of polyamines, molecules strongly associated with many biological processes including protection from osmotic stress [60]. TCA cycle-derived amino acids of the glutamate and aspartate families are central regulators of carbon and nitrogen metabolism [61]. In fact, a high level of glutamine synthetase activity has been found in the leaves of *C. arabica* cv. Catual during grain expansion [62], which has been linked to the capacity of coffee plants to assimilate nitrogen, a useful trait to improve coffee management in commercial orchards.

In Icatu, amino acids derived from the shikimate pathway (e.g., phenylalanine, tryptophan, and tyrosine) accumulated under single exposure to SWD under aCO<sub>2</sub>; tryptophan and tyrosine also increased in SWD under eCO<sub>2</sub> conditions. The accumulation of these metabolites is a valuable source of carbon-skeletons for the phenylpropanoid pathway and further biosynthesis of secondary metabolites [63,64]. Tryptophan is reported to play an important role as an osmolyte, but also in stomatal regulation and reactive oxygen species (ROS) scavenging under drought conditions [65].

A significant increase in putrescine occurred in Icatu under the interaction of SWD and  $eCO_2$ . Several drought experiments have shown a correlation between resistance to dehydration and the levels of putrescine—a plant growth regulator that helps to balance chlorophyll concentrations—and the accumulation of osmolytes and soluble phenolic compounds under drought conditions [66].

Osmoprotective sugars were also affected by drought and a significant increase was observed in fructose in Icatu and in glucose, trehalose, and sucrose in CL153 (particularly under SWD and eCO<sub>2</sub> conditions), which might have contributed to dehydration

tolerance [36,37]. The metabolite responses of each genotype to water deficit and eCO<sub>2</sub>, together with the PCA and PLS-DA analysis, showed that eCO<sub>2</sub> clearly altered the impact of MWD in both genotypes, as shown by the overlap of the MWD under eCO<sub>2</sub> cluster with WW controls. However, MWD plants under aCO<sub>2</sub> did not show such proximity to WW controls. This observation is in line with the positive effect of eCO<sub>2</sub> regarding the water deficit imposition reported earlier [28,32], mainly in Icatu plants. In addition, the clear separation of SWD plants from the WW and MWD clusters, particularly in Icatu plants, showed that the production of metabolites was greatly altered by SWD. Nevertheless, Icatu plants maintained (or increased) their metabolite levels under both CO<sub>2</sub> concentrations (within MWD and SWD), contrasting with the decreases observed in CL153 under eCO<sub>2</sub>, which is in line with the high tolerance of Icatu plants to water deficit under both CO<sub>2</sub> concentrations [32].

Physiological and biochemical studies demonstrated an absence of impact on Icatu plants in most parameters under SWD, irrespective of  $CO_2$  concentration, and a greater tolerance than CL153 under aCO<sub>2</sub>. This resilience of Icatu was associated with the preservation of photosynthetic performance (e.g., photosynthetic capacity,  $A_{max}$ ; photochemical efficiency of PSII,  $F_v/F_m$ ) and the maintenance or reinforcement of several chloroplast components (e.g., cytochromes electron carriers, photosystems (PSs), and RuBisCO activities) [32,44]. Such intrinsic tolerance (reinforced under eCO<sub>2</sub>) was further associated with a wide abundance of proteins related to the photosynthetic apparatus (as part of photosystems, light harvesting complexes, protective cyclic electron flow, and RuBisCO activase), which was not observed in CL153 [32].

Limited photochemical energy use can generate ROS leading to damage to the photosynthetic apparatus (e.g., D1 protein, lipids, electron transport) [67,68]. Therefore, at the cellular level, drought tolerance is often associated with the activation of photoprotective and antioxidative mechanisms. This has previously been shown to be a key response in the acclimation of coffee plants to a wide variety of abiotic stresses, such as high irradiance [69,70], cold [71–73], and temperature [19,21,31]. Antioxidant enzymes coupled with strong lipid dynamics usually contribute to the maintenance of photosynthetic activity in coffee genotypes (including in Icatu). For instance, drought experiments using *C. canephora* plants of Clone 120 (drought-tolerant) and Clone 109 (drought-sensitive), subjected to repetitive cycles of drought conditions for 14 days, revealed an oxidative stress response on the sensitive clone, presumably leading to programmed cell death, while acclimation of the tolerant genotype seems to be related to antioxidant secondary metabolites and ABA response [74].

#### 4. Materials and Methods

#### 4.1. Plant Material and Growth Conditions

Plants of two cropped genotypes (Brazil) from the two main coffee-producing species, *Coffea canephora* Pierre ex A. Froehner cv. Conilon Clone 153 (CL153) and *C. arabica* L. cv. Icatu Vermelho (an introgressed cultivar resulting from a cross of *C. canephora* and *C. arabica* cv. Bourbon Vermelho, which was then further crossed with *C. arabica* cv. Mundo Novo) were used, as described in [32]. Briefly, plants were grown for seven years in 80 L pots in two walk-in growth chambers (EHHF 10000, ARALAB, Portugal), under controlled temperature conditions (25/20 °C, day/night,  $\pm 1$  °C), irradiance (ca. 750 µmol m<sup>-2</sup> s<sup>-1</sup> at the upper part of the plant), relative humidity (70%  $\pm$  2%), photoperiod (12 h), and either ambient (aCO<sub>2</sub>, 380  $\pm$  5 ppm) or elevated (eCO<sub>2</sub>, 700  $\pm$  5 ppm) air CO<sub>2</sub> concentration. Plants were grown without restriction of nutrients [14] or root growth, and with adequate water availability provided by watering the plants every two days.

#### 4.2. Water Deficit Imposition

Within each chamber, plants were divided into three groups (n = 6 to 8) and maintained under well-irrigated conditions (WW), or exposed to moderate (MWD) and severe water deficit (SWD), as described in [32]. Water deficit was gradually imposed over two weeks by partially withholding irrigation (that is, by replacing only part of the amount of lost water per pot). Water availability levels were established as a function of leaf water potential at predawn ( $\Psi_{pd}$ ), as WW:  $\Psi_{pd} > -0.35$  MPa; MWD  $-1.5 < \Psi_{pd} < 2.5$  MPa; and SWD  $\Psi_{pd} < 3.5$  MPa, corresponding to ca. 80% (WW), 25% (MWD), or 10% (SWD) of maximal pot water availability, respectively [75].  $\Psi_{pd}$  was determined immediately after leaf excision in 5–6 plants per treatment, using a pressure chamber (Model 1000, PMS Instrument Co., Albany, OR, USA).

When the required  $\Psi_{pd}$  was reached for MWD or SWD conditions, pot moisture was maintained for another two weeks by adding adequate water amounts according to each watering treatment before sampling. Exceptionally, the Icatu 700-plants (eCO<sub>2</sub>) under MWD conditions were exposed to total water withholding in the last five days of the four-week period, in order to further force the reduction of  $\Psi_{pd}$  values, which did not shift below -0.6 MPa until that stage.

For metabolomic analysis, newly matured leaves from the upper third part (well illuminated) were collected, immediately frozen in liquid nitrogen, and stored at -80 °C.

## 4.3. GC-TOF-MS Primary Metabolite Profiling

Primary metabolites were extracted and derivatized following a well-established protocol [76]. Frozen leaves were ground to a fine powder with a mortar and pestle and under liquid nitrogen. A total of 50 mg fresh weight (FW) of finely homogenized leaf material was weighed into each 1.5 mL safe-lock polypropylene microfuge tube and 700  $\mu$ L of ice-cold methanol and 30  $\mu$ L of ribitol (0.2 mg mL<sup>-1</sup> ribitol in water), as the internal standard, were added to each tube. Samples were vortex-mixed and incubated in a shaker (ThermoMixer, Eppendorf, Hamburg, Germany) for 15 min at 70 °C and 950 rpm, and subsequently centrifuged ( $12,000 \times g$ , 10 min, 20 °C). The supernatant was collected, mixed with 375 µL chloroform and 750 µL water, and vortex-mixed. After centrifugation  $(2200 \times g, 15 \text{ min}, 20 \degree \text{C}), 150 \text{ mL}$  of the polar aqueous/methanol phase was evaporated to dryness using a centrifugal concentrator for 3 h at 30 °C (Vacufuge Plus, Eppendorf, Hamburg, Germany), and stored at -80 °C. Dried polar extracts were derivatized with 40  $\mu$ L of 20 mg mL<sup>-1</sup> methoxyamine hydrochloride in pyridine, followed by 70  $\mu$ L of *N*-methyl-*N*-trimethylsilyltrifluoroacetamide (TMS derivatization) and 20  $\mu$ L mL<sup>-1</sup> of a mixture of fatty acid methyl esters (FAMES). Biological variations were controlled by analyzing quality control (QC) standards; namely, FAMES internal standard markers and a QC standard solution of 41 pure reference compounds, throughout the analysis. After GC-TOF-MS analysis, samples were subsequently evaluated using AMDIS (Automated Mass Spectral Deconvolution and Identification System software, version 2.71). Primary metabolites were annotated using the TagFinder software [77] and a reference library of ambient mass spectra and retention indices from the Golm Metabolome Database (http://gmd.mpimp-golm.mpg.de/) [78,79]. Metadata information following minimum reporting standard guidelines of the Metabolomics Standard Initiative (MSI) can be found in Supplementary Table S3.

#### 4.4. Experimental Design and Statistical Analyses

Plants from each coffee genotype were subjected to six treatment combinations, reflecting the single and combined exposure to two  $CO_2$  concentration (a $CO_2$  or e $CO_2$ ) and three levels of water availability (WW, MWD or SWD). The experiment was implemented using a completely randomized design, with six to eight plants per treatment in individual pots.

Statistical analyses were performed in R and R Studio software [48,49]. R Packages used to perform statistical analysis include "agricolae" [80], "gplots" [50], and "mixOmics" [51].

The relative abundance of primary metabolite levels was normalized to the internal standard (ribitol) and the fresh weight/dry weight ratio of the samples. Within each genotype, one-way ANOVA at a 95% confidence level was used to assess differences between treatments. Subsequently, fold-changes between conditions and the control for each metabolite were determined and  $Log_{10}$  transformed for heatmap plotting.

Principal component analysis (PCA) was applied using unit variance scaling, and to further maximize the discrimination between the sample groups, supervised partial least squares discriminant analysis (PLS-DA) was performed using the leave-one-out cross-validation embedded in the "mixOmics" package.

## 5. Conclusions

Overall, our metabolite profiling findings showed significant drought impacts on two widely traded coffee species. Notably,  $eCO_2$  clearly reduced metabolite changes under MWD in Icatu (but not in CL153), resulting in metabolic profiles close to those of their WW plants. These findings are relevant in terms of the coffee plant response to future climate change scenarios (particularly in regard to the role of  $eCO_2$ ), and also highlight the need for complementary and integrative studies, from ecophysiology to omics, to better clarify the role of specific metabolites in the resilience of this important tropical crop.

**Supplementary Materials:** The following are available online at https://www.mdpi.com/article/ 10.3390/metabo11070427/s1, Table S1: Fold changes in relative levels of primary metabolites in leaves of *Coffea canephora* cv. Conilon: Clone 153, Table S2: Fold changes in relative levels of primary metabolites in leaves of *Coffea arabica* cv. Icatu, Table S3: Metabolomics Standards Initiative compliant metadata.

Author Contributions: Conceptualization, A.I.R.-B., F.C.L., F.M.D., I.M. and J.C.R.; methodology, A.I.R.-B., C.A. and J.C.R.; software, A.M.R.; validation, C.A.; formal analysis, A.M.R.; investigation, A.M.R., D.M.P., J.C.R. and T.J.; resources, C.A., J.C.R. and S.O.; data curation, A.M.R.; writing—original draft preparation, A.M.R.; writing—review and editing, A.I.R.-B., C.A., D.M.P., F.C.L., F.M.D., I.M., J.C.R., S.O. and T.J.; visualization, A.M.R.; supervision, C.A.; project administration, A.I.R.-B. and J.C.R.; funding acquisition, A.I.R.-B. and J.C.R. All authors have read and agreed to the published version of the manuscript.

**Funding:** This work received funding support from the European Union's Horizon 2020 research and innovation program (grant agreement No. 727934, project BreedCAFS), and from national funds from Fundação para a Ciência e a Tecnologia (FCT), Portugal, through the project PTDC/ASP-AGR/31257/2017, and the research units UIDB/00239/2020 (CEF), UIDP/04035/2020 (GeoBioTec), and UID/Multi/04551/2013 (GREEN-IT). C.A. acknowledges the FCT Investigator Programme (contract IF/00376/2012/CP0165/CT0003). A.M.R. acknowledges FCT and the ITQB NOVA international Ph.D. programme 'Plants for Life' (PD/00035/2013) for the Ph.D. fellowship (PD/BD/114417/2016). T.J. acknowledges FCT and the ITQB NOVA international Ph.D. fellowship (PD/BD/113475/2015). Fellowships from the Conselho Nacional de Desenvolvimento Científico e Tecnológico, Brazil (CNPq) and Fundação de Amparo à Pesquisa do Estado de Minas Gerais, Brazil (FAPEMIG, project CRA-RED-00053-16) to F.M.D. are also gratefully acknowledged.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

**Data Availability Statement:** The data presented in this study are available in article and Supplementary Materials.

**Conflicts of Interest:** The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; or in the decision to publish the results.

#### References

- 1. FAO Statistical Pocketbook; Coffee Food and Agricultural Organization of the United Nations: Rome, Italy, 2015.
- 2. DaMatta, F.M.; Rahn, E.; Läderach, P.; Ghini, R.; Ramalho, J.C. Why could the coffee crop endure climate change and global warming to a greater extent than previously estimated? *Clim. Chang.* **2019**, *152*, 167–178. [CrossRef]

- Ramirez-Villegas, J.; Challinor, A. Assessing relevant climate data for agricultural applications. *Agric. For. Meteorol.* 2012, 161, 26–45. [CrossRef]
- 4. Magrach, A.; Ghazoul, J. Climate and pest-driven geographic shifts in global coffee production: Implications for forest cover, biodiversity and carbon storage. *PLoS ONE* **2015**, *10*, e0133071. [CrossRef] [PubMed]
- Ovalle-Rivera, O.; L\u00e4derach, P.; Bunn, C.; Obersteiner, M.; Schroth, G. Projected shifts in *Coffea arabica* suitability among major global producing regions due to climate change. *PLoS ONE* 2015, *10*, e0124155. [CrossRef]
- 6. van der Vossen, H.; Bertrand, B.; Charrier, A. Next generation variety development for sustainable production of arabica coffee (*Coffea arabica* L.): A review. *Euphytica* 2015, 204, 243–256. [CrossRef]
- Läderach, P.; Ramirez–Villegas, J.; Navarro-Racines, C.; Zelaya, C.; Martinez-Valle, A.; Jarvis, A. Climate change adaptation of coffee production in space and time. *Clim. Chang.* 2017, 141, 47–62. [CrossRef]
- 8. Davis, A.P.; Chadburn, H.; Moat, J.; O'Sullivan, R.; Hargreaves, S.; Lughadha, E.N. High extinction risk for wild coffee species and implications for coffee sector sustainability. *Sci. Adv.* **2019**, *5*, eaav3473. [CrossRef] [PubMed]
- 9. de Sousa, K.; van Zonneveld, M.; Holmgren, M.; Kindt, R.; Ordoñez, J.C. The future of coffee and cocoa agroforestry in a warmer Mesoamerica. *Sci. Rep.* **2019**, *9*, 8828. [CrossRef]
- 10. IPCC Core Writing Team. Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change; IPCC: Geneva, Switzerland, 2014.
- 11. DaMatta, F.M.; Grandis, A.; Arenque, B.C.; Buckeridge, M.S. Impacts of climate changes on crop physiology and food quality. *Food Res. Int.* **2010**, *43*, 1814–1823. [CrossRef]
- 12. Ainsworth, E.A.; Rogers, A. The response of photosynthesis and stomatal conductance to rising [CO<sub>2</sub>]: Mechanisms and environmental interactions. *Plant Cell Environ.* **2007**, *30*, 258–270. [CrossRef] [PubMed]
- 13. Kirschbaum, M.U.F. Does enhanced photosynthesis enhance growth? Lessons learned from CO<sub>2</sub> enrichment studies. *Plant Physiol.* **2011**, *155*, 117–124. [CrossRef] [PubMed]
- Ramalho, J.C.; Rodrigues, A.P.; Semedo, J.N.; Pais, I.P.; Martins, L.D.; Simões-Costa, M.C.; Leitão, A.E.; Fortunato, A.S.; Batista-Santos, P.; Palos, I.M.; et al. Sustained photosynthetic performance of *Coffea* spp. under long-term enhanced [CO<sub>2</sub>]. *PLoS ONE* 2013, *8*, e82712. [CrossRef] [PubMed]
- 15. Ghini, R.; Torre-Neto, A.; Dentzien, A.F.M.; Guerreiro-Filho, O.; Lost, R.; Patrício, F.R.A.; Prado, J.S.M.; Thomaziello, R.A.; Bettiol, W.; DaMatta, F.M. Coffee growth, pest and yield responses to free-air CO<sub>2</sub> enrichment. *Clim. Chang.* **2015**, *132*, 307–320. [CrossRef]
- DaMatta, F.M.; Godoy, A.G.; Menezes-Silva, P.E.; Martins, S.C.V.; Sanglard, L.M.V.P.; Morais, L.E.; Torre-Neto, A.; Ghini, R. Sustained enhancement of photosynthesis in coffee trees grown under free-air CO<sub>2</sub> enrichment conditions: Disentangling the contributions of stomatal, mesophyll, and biochemical limitations. *J. Exp. Bot.* 2016, *67*, 341–352. [CrossRef]
- 17. DaMatta, F.M.; Avila, R.T.; Cardoso, A.A.; Martins, S.C.V.; Ramalho, J.C. Physiological and agronomic performance of the coffee crop in the context of climate change and global warming: A review. J. Agric. Food Chem. 2018, 66, 5264–5274. [CrossRef]
- Rakocevic, M.; Braga, K.S.M.; Batista, E.R.; Maia, A.H.N.; Scholz, M.B.S.; Filizola, H.F. The vegetative growth assists to reproductive responses of Arabic coffee trees in a long-term FACE experiment. *Plant Growth Reg.* 2020, 91, 305–316. [CrossRef]
- Martins, M.Q.; Rodrigues, W.P.; Fortunato, A.S.; Leitão, A.E.; Rodrigues, A.P.; Pais, I.P.; Martins, L.D.; Silva, M.J.; Reboredo, F.H.; Partelli, F.L.; et al. Protective response mechanisms to heat stress in interaction with high [CO<sub>2</sub>] conditions in *Coffea* spp. *Front. Plant Sci.* 2016, 7, 947. [CrossRef]
- Rodrigues, W.P.; Martins, M.Q.; Fortunato, A.S.; Rodrigues, A.P.; Semedo, J.N.; Simoes-Costa, M.C.; Pais, I.P.; Leitão, A.E.; Colwell, F.; Goulão, L.; et al. Long-term elevated air [CO<sub>2</sub>] strengthens photosynthetic functioning and mitigates the impact of supra-optimal temperatures in tropical *Coffea arabica* and *Coffea canephora* species. *Glob. Chang. Biol.* 2016, 22, 415–431. [CrossRef]
- Scotti-Campos, P.; Pais, I.P.; Ribeiro-Barros, A.I.; Martins, L.D.; Tomaz, M.A.; Rodrigues, W.P.; Campostrini, E.; Semedo, J.N.; Fortunato, A.S.; Martins, M.Q.; et al. Lipid profile adjustments may contribute to warming acclimation and to heat impact mitigation by elevated [CO<sub>2</sub>] in *Coffea* spp. *Environ. Exp. Bot.* 2019, *167*, 103856. [CrossRef]
- 22. Martins, L.D.; Tomaz, M.A.; Lidon, F.C.; DaMatta, F.M.; Ramalho, J.C. Combined effects of elevated [CO<sub>2</sub>] and high temperature on leaf mineral balance in *Coffea* spp. plants. *Clim. Chang.* **2014**, *126*, 365–379. [CrossRef]
- Ramalho, J.C.; Pais, I.P.; Leitão, A.E.; Guerra, M.; Reboredo, F.H.; Máguas, C.M.; Carvalho, M.L.; Scotti-Campos, P.; Ribeiro-Barros, A.I.; Lidon, F.C.; et al. Can elevated air [CO<sub>2</sub>] conditions mitigate the predicted warming impact on the quality of coffee bean? *Front. Plant Sci.* 2018, 9, 287. [CrossRef]
- 24. Tausz-Posch, S.; Tausz, M.; Bourgault, M. Elevated [CO<sub>2</sub>] effects on crops: Advances in understanding acclimation, nitrogen dynamics and interactions with drought and other organisms. *Plant Biol.* **2020**, *22*, 38–51. [CrossRef] [PubMed]
- Koutavas, A. CO<sub>2</sub> fertilization and enhanced drought resistance in Greek firs from Cephalonia Island, Greece. *Glob. Chang. Biol.* 2013, 19, 529–539. [CrossRef]
- Wang, A.; Lam, S.K.; Hao, X.; Li, F.Y.; Zong, Y.; Wang, H.; Li, P. Elevated CO<sub>2</sub> reduces the adverse effects of drought stress on a high-yielding soybean (*Glycine max* (L.) Merr.) cultivar by increasing water use efficiency. *Plant Physiol. Biochem.* 2018, 132, 660–665. [CrossRef] [PubMed]
- Gamar, M.I.A.; Kisiala, A.; Emery, R.J.N.; Yeung, E.C.; Stone, S.L.; Qaderi, M.M. Elevated carbon dioxide decreases the adverse effects of higher temperature and drought stress by mitigating oxidative stress and improving water status in *Arabidopsis thaliana*. *Planta* 2019, 250, 1191–1214. [CrossRef]

- Avila, R.T.; Almeida, W.L.; Costa, L.C.; Machado, K.L.; Barbosa, M.L.; de Souza, R.P.; Martino, P.B.; Juárez, M.A.T.; Marçal, D.M.S.; Martins, S.C.V.; et al. Elevated air [CO<sub>2</sub>] improves photosynthetic performance and alters biomass accumulation and partitioning in drought-stressed coffee plants. *Environ. Exp. Bot.* 2020, 177, 104137. [CrossRef]
- Avila, R.T.; Cardoso, A.A.; de Almeida, W.L.; Costa, L.C.; Machado, K.L.G.; Barbosa, M.L.; de Souza, R.P.B.; Oliveira, L.A.; Batista, D.S.; Martins, S.C.V.; et al. Coffee plants respond to drought and elevated [CO<sub>2</sub>] through changes in stomatal function, plant hydraulic conductance, and aquaporin expression. *Environ. Exp. Bot.* 2020, *177*, 104148. [CrossRef]
- Marques, I.; Fernandes, I.; David, P.H.C.; Paulo, O.S.; Goulao, L.F.; Fortunato, A.S.; Lidon, F.C.; DaMatta, F.M.; Ramalho, J.C.; Ribeiro-Barros, A.I. Transcriptomic leaf profiling reveals differential responses of the two most traded coffee species to elevated [CO<sub>2</sub>]. *Int. J. Mol. Sci.* 2020, *21*, 9211. [CrossRef]
- Marques, I.; Fernandes, I.; Paulo, O.S.; Lidon, F.C.; DaMatta, F.M.; Ramalho, J.C.; Ribeiro-Barros, A.I. A transcriptomic approach to understanding the combined impacts of supra-optimal temperatures and CO<sub>2</sub> revealed different responses in the polyploid *Coffea arabica* and its diploid progenitor *C. canephora. Int. J. Mol. Sci.* 2021, 22, 3125. [CrossRef] [PubMed]
- Semedo, J.N.; Rodrigues, A.P.; Lidon, F.C.; Pais, I.P.; Marques, I.; Gouveia, D.; Armengaud, J.; Silva, M.J.; Martins, S.; Semedo, M.C.; et al. Intrinsic non-stomatal resilience to drought of the photosynthetic apparatus in *Coffea* spp. is strengthened by elevated air [CO<sub>2</sub>]. *Tree Physiol.* 2021, 41, 708–727. [CrossRef]
- 33. Mittler, R. Abiotic stress, the field environment and stress combination. Trends Plant Sci. 2006, 11, 15–19. [CrossRef] [PubMed]
- 34. Mittler, R.; Blumwald, E. Genetic engineering for modern agriculture: Challenges and perspectives. *Annu. Rev. Plant Biol.* **2010**, 61, 443–462. [CrossRef]
- Suzuki, N.; Rivero, R.M.; Shulaev, V.; Blumwald, E.; Mittler, R. Abiotic and biotic stress combinations. New Phytol. 2014, 203, 32–43. [CrossRef] [PubMed]
- 36. Obata, T.; Fernie, A.R. The use of metabolomics to dissect plant responses to abiotic stresses. *Cell. Mol. Life Sci.* 2012, *69*, 3225–3243. [CrossRef]
- Jorge, T.F.; António, C. Plant Metabolomics in a Changing World: Metabolite Responses to Abiotic Stress Combinations, Plant, Abiotic Stress and Responses to Climate Change. In *Plant, Abiotic Stress and Responses to Climate Change*; Andjelkovic, V., Ed.; InTechOpen Publishing: London, UK, 2017; pp. 111–132. [CrossRef]
- Chaves, M.M.; Oliveira, M.M. Mechanisms underlying plant resilience to water deficits: Prospects for water-saving agriculture. J. Exp. Bot. 2004, 55, 2365–2384. [CrossRef]
- 39. Fahad, S.; Bajwa, A.A.; Nazir, U.; Anjum, S.A.; Farooq, A.; Zohaib, A.; Sadia, S.; Nasim, W.; Adkins, S.; Saud, S.; et al. Crop production under drought and heat stress: Plant responses and management options. *Front. Plant Sci.* **2017**, *8*, 1147. [CrossRef]
- 40. Hare, P.D.; Cress, W.A.; van Staden, J. Dissecting the roles of osmolyte accumulation during stress. *Plant Cell Environ*. **1998**, 21, 535–553. [CrossRef]
- 41. Krasensky, J.; Jonak, C. Drought, salt and temperature stress induced metabolic rearrangements and regulatory networks. *J. Exp. Bot.* 2012, *63*, 1593–1608. [CrossRef]
- 42. Arbona, V.; Manzi, M.; de Ollas, C.; Gómez-Cadenas, A. Metabolomics as a tool to investigate abiotic stress tolerance in plants. *Int. J. Mol. Sci.* 2013, 14, 4885–4911. [CrossRef]
- Jorge, T.F.; Rodrigues, J.A.; Caldana, C.; Sçhimt, R.; van Dongen, J.; Thomas-Oates, J.; António, C. Mass spectrometry-based plant metabolomics: Metabolite responses to abiotic stress. *Mass Spec. Rev.* 2016, 35, 620–649. [CrossRef] [PubMed]
- Dubberstein, D.; Lidon, F.C.; Rodrigues, A.P.; Semedo, J.N.; Marques, I.; Rodrigues, W.P.; Gouveia, D.; Armengaud, J.; Semedo, M.C.; Martins, S.; et al. Resilient and sensitive key points of the photosynthetic machinery of *Coffea* spp. to the single and superimposed exposure to severe drought and heat stresses. *Front. Plant Sci.* 2020, *11*, 1049. [CrossRef]
- Fiehn, O.; Kopka, J.; Dörmann, P.; Tretheway, R.N.; Willmitzer, L. Metabolite profiling of plant functional genomics. *Nat. Biotechnol.* 2000, 18, 1157–1161. [CrossRef]
- 46. Fernie, A. Metabolome characterisation in plant system analysis. Funct. Plant Biol. 2003, 30, 111–120. [CrossRef] [PubMed]
- 47. Kopka, J.; Fernie, A.; Weckwerth, W.; Gibon, Y.; Stitt, M. Metabolite profiling in plant biology: Platforms and destinations. *Genome Biol.* **2004**, *5*, 109. [CrossRef]
- 48. R Core Team. *R: A Language and Environment for Statistical Computing;* R Foundation for Statistical Computing: Vienna, Austria, 2020. Available online: https://www.R-project.org/ (accessed on 24 November 2020).
- 49. RStudio Team. *RStudio: Integrated development for R*; RStudio, Inc.: Boston, MA, USA, 2020; Available online: http://www.rstudio. com/ (accessed on 24 November 2020).
- Warnes, G.R.; Bolker, B.; Bonebakker, L.; Gentleman, R.; Huber, W.; Liaw, A.; Thomas, L.; Maechler, M.; Magnusson, A.; Moeller, S.; et al. gplots: Various R Programming Tools for Plotting Data; R Package Version 3.0.3; 2020. Available online: https://CRAN.R-project.org/package=gplots (accessed on 24 November 2020).
- 51. Rohart, F.; Gautier, B.; Singh, A.; Lê Cao, K.-A. mixOmics: An R package for 'omics feature selection and multiple data integration. *PLoS Comput. Biol.* **2017**, *13*, e1005752. [CrossRef]
- Mata, A.T.; Jorge, T.F.; Pires, M.V.; António, C. Drought stress tolerance in plants: Insights from metabolomics. In *Drought Stress Tolerance in Plants, Molecular and Genetic Perspectives*; Hossain, M.A., Wani, S.H., Bhattachajee, S., Burritt, D.J., Phan, L.S., Eds.; Tran. Humana Press, Springer International Publishing: Cham, Switzerland, 2016; Volume 2, pp. 187–216. [CrossRef]

- 53. Persson, T.; Van Nguyen, T.; Alloisio, N.; Pujic, P.; Berry, A.M.; Normand, P.; Pawlowski, K. The N-metabolites of roots and actinorhizal nodules from *Alnus glutinosa* and *Datisca glomerata*: Can *D. glomerata* change N-transport forms when nodulated? *Symbiosis* 2016, 70, 149–157. [CrossRef]
- 54. Araújo, W.L.; Ishizaki, K.; Nunes-Nesi, A.; Larson, T.R.; Tohge, T.; Krahnert, I.; Witt, S.; Obata, T.; Schauer, N.; Graham, I.A.; et al. Identification of the 2-hydroxyglutarate and isovaleryl-CoA dehydrogenases as alternative electron donors linking lysine catabolism to the electron transport chain of Arabidopsis mitochondria. *Plant Cell* **2010**, *22*, 1549–1563. [CrossRef]
- 55. Pires, M.V.; Pereira Júnior, A.A.; Medeiros, D.B.; Daloso, D.M.; Pham, P.A.; Barros, K.A.; Engqvist, M.K.; Florian, A.; Krahnert, I.; Maurino, V.G.; et al. The influence of alternative pathways of respiration that utilize branched-chain amino acids following water shortage in Arabidopsis. *Plant Cell Environ.* **2016**, *39*, 1304–1319. [CrossRef] [PubMed]
- 56. Tamirat, W. Review on role of proline on coffee under drought conditions. J. Environ. Earth Sci. 2019, 9, 10. [CrossRef]
- Silva, V.A.; Antunes, W.C.; Guimarães, B.L.S.; Paiva, R.M.C.; Silva, V.F.; Ferrão, M.A.G.; DaMatta, F.M.; Loureiro, M.E. Physiological response of Conilon coffee clone sensitive to drought grafted onto tolerant rootstock. *Pesq. Agropec. Bras.* 2010, 45, 457–464. [CrossRef]
- 58. Santos, A.B.; Mazzafera, P. Dehydrins are highly expressed in water-stressed plants of two coffee species. *Trop. Plant Biol.* **2012**, *5*, 218–232. [CrossRef]
- Kalamaki, M.S.; Alexandrou, D.; Lazari, D.; Merkouropoulos, G.; Fotopoulos, V.; Pateraki, I.; Aggelis, A.; Carrillo-López, A.; Rubio-Cabetas, M.J.; Kanellis, A.K. Over-expression of a tomato N-acetyl-L-glutamate synthase gene (SINAGS1) in *Arabidopsis thaliana* results in high ornithine levels and increased tolerance in salt and drought stresses. J. Exp. Bot. 2009, 60, 1859–1871. [CrossRef]
- Kalamaki, M.S.; Merkouropoulos, G.; Kanellis, A.K. Can ornithine accumulation modulate abiotic stress tolerance in Arabidopsis? *Plant Signal. Behav.* 2009, 4, 1099–1101. [CrossRef] [PubMed]
- 61. Fait, A.; Fromm, H.; Walter, D.; Galili, G.; Fernie, A.R. Highway or byway: The metabolic role of the GABA shunt in plants. *Trends Plant Sci.* **2008**, *13*, 14–19. [CrossRef]
- 62. Neto, A.P.; Favarin, J.L.; dos Reis, A.R.; Tezzoto, T.; de Almeida, R.E.M.; Junior, J.L.; Fallo, L.A. Nitrogen metabolism in coffee plants in response to nitrogen supply by fertigation. *Theor. Exp. Plant Physiol.* **2015**, 27, 41–50. [CrossRef]
- 63. Tohge, T.; Watanabe, M.; Hoefgen, R.; Fernie, A.R. Shikimate and phenylalanine biosynthesis in the green lineage. *Front. Plant Sci.* **2013**, *4*, 62. [CrossRef] [PubMed]
- 64. Hildebrandt, T.M. Synthesis versus degradation: Directions of amino acid metabolism during Arabidopsis abiotic stress response. *Plant Mol. Biol.* **2018**, *98*, 121–135. [CrossRef]
- 65. Bowne, J.B.; Erwin, T.A.; Juttner, J.; Schnurbusch, T.; Langridge, P.; Bacic, A.; Roessner, U. Drought responses of leaf tissues from wheat cultivars of differing drought tolerance at the metabolite level. *Mol. Plant* **2012**, *5*, 418–429. [CrossRef]
- Alcázar, R.; Planas, J.; Saxena, T.; Zarza, X.; Bortolotti, C.; Cuevas, J.; Bitrián, M.; Tiburcio, A.F.; Altabella, T. Putrescine accumulation confers drought tolerance in transgenic Arabidopsis plants over-expressing the homologous Arginine decarboxylase 2 gene. *Plant Physiol. Biochem.* 2010, 48, 547–552. [CrossRef]
- 67. Chaves, M.M.; Flexas, J.; Pinheiro, C. Photosynthesis under drought and salt stress: Regulation mechanisms from whole plant to cell. *Ann. Bot.* **2009**, *103*, 551–560. [CrossRef] [PubMed]
- 68. Osakabe, Y.; Osakabe, K.; Shinozaki, K.; Tran, L.S.P. Response of plants to water stress. Front. Plant Sci. 2014, 5, 86. [CrossRef]
- Ramalho, J.C.; Campos, P.S.; Teixeira, M.; Nunes, M.A. Nitrogen dependent changes in antioxidant systems and in fatty acid composition of chloroplast membranes from *Coffea arabica* L. plants submitted to high irradiance. *Plant Sci.* 1998, 135, 115–124. [CrossRef]
- Ramalho, J.; Campos, P.; Quartin, V.; Silva, M.J.; Nunes, M.A. High irradiance impairments on photosynthetic electron transport, ribulose-1,5-bisphosphate carboxylase/ oxygenase and N assimilation as a function of N availability in *Coffea arabica* L. plants. J. *Plant Physiol.* 1999, 154, 319–326. [CrossRef]
- Fortunato, A.; Lidon, F.; Eichler, P.; Leitão, A.; Pais, I.; Ribeiro, A.; Ramalho, J. Biochemical and molecular characterization of the antioxidative system of *Coffea* sp. under cold conditions in genotypes with contrasting tolerance. *J. Plant Physiol.* 2010, 167, 333–342. [CrossRef]
- 72. Scotti-Campos, P.; Pais, I.P.; Partelli, F.L.; Batista-Santos, P.; Ramalho, J.C. Phospholipids profile in chloroplasts of *Coffea* spp. genotypes differing in cold acclimation ability. *J. Plant Physiol.* **2014**, 171, 243–249. [CrossRef]
- 73. Ramalho, J.C.; DaMatta, F.M.; Rodrigues, A.P.; Scotti-Campos, P.; Pais, I.; Batista-Santos, P.; Partelli, F.L.; Ribeiro, A.; Lidon, F.C.; Leitão, A.E. Cold impact and acclimation response of *Coffea* spp. plants. *Theor. Exp. Plant Physiol.* **2014**, *26*, 5–18. [CrossRef]
- 74. Guedes, F.A.F.; Nobres, P.; Ferreira, D.C.R.; Menezes-Silva, P.E.; Ribeiro-Alves, M.; Correa, R.L.; DaMatta, F.M.; Alves-Ferreira, M. Transcriptional memory contributes to drought tolerance in coffee (*Coffea canephora*) plants. *Environ. Exp. Bot.* 2018, 147, 220–233. [CrossRef]
- 75. Ramalho, J.C.; Rodrigues, A.P.; Lidon, F.C.; Marques, L.M.C.; Leitão, A.E.; Fortunato, A.F.; Pais, I.P.; Silva, M.J.; Scotti-Campos, P.; Lopes, A.; et al. Stress cross-response of the antioxidative system promoted by superimposed drought and cold conditions in *Coffea* spp. *PLoS ONE* **2018**, *13*, e0198694. [CrossRef]
- 76. Lisec, J.; Schauer, N.; Kopka, J.; Willmitzer, L.; Fernie, A.R. Gas chromatography mass spectrometry–based metabolite profiling in plants. *Nat. Protoc.* 2006, *1*, 387–396. [CrossRef]

- 77. Luedemann, A.; Strassburg, K.; Erban, A.; Kopka, J. TagFinder for the quantitative analysis of gas chromatography-mass spectrometry (GC-MS)-based metabolite profiling experiments. *Bioinformatics* **2008**, *24*, 732–737. [CrossRef]
- 78. Kopka, J.; Schauer, N.; Krueger, S.; Birkemeyer, C.; Usadel, B.; Bergmüller, E.; Dörmann, P.; Weckwerth, W.; Gibon, Y.; Stitt, M.; et al. GMD@CSB.DB: The golm metabolome database. *Bioinformatics* **2005**, *21*, 1635–1638. [CrossRef] [PubMed]
- Schauer, N.; Steinhauser, D.; Strelkov, S.; Schomburg, D.; Allison, G.; Moritz, T.; Lundgren, K.; Roessner-Tunali, U.; Forbes, M.G.; Willmitzer, L.; et al. GC-MS libraries for the rapid identification of metabolites in complex biological samples. *FEBS Lett.* 2005, 579, 1332–1337. [CrossRef] [PubMed]
- 80. Mendiburu, F. *Agricolae: Statistical Procedures for Agricultural Research;* R Package Version 1.3-3; 2020. Available online: https://CRAN.R-project.org/package=agricolae (accessed on 24 November 2020).